

# Cooking well

Delicious, everyday  
recipes for the PKD family

Jacob Taylor, Ph.D., RD, LD



**PKD FOUNDATION**  
Polycystic Kidney Disease



# Cooking **well**

Delicious,  
everyday recipes  
for the PKD family



**PKD FOUNDATION**  
Polycystic Kidney Disease

## Cooking Well

*Delicious, everyday recipes for the PKD family*

Copyright © 2016 PKD Foundation.

All rights reserved. No part of this book may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording or by any information storage and retrieval system without prior written permission from the publisher.

ISBN: 978-0-692-83054-3

### ACKNOWLEDGMENTS

#### AUTHOR

**Jacob Taylor, Ph.D., RD, LD**

#### CONTRIBUTORS

**Osama Amro, M.D.**

Swedish Medical Center, Seattle, WA

Provided content for the water and ADPKD section.

**Kelly L. Welsh, RD, CD**

Author of "Brilliant Eats," the original kidney-friendly cookbook from which some of the recipes in this book were taken.

Recipe analysis was conducted at the University of Kansas Medical Center Department of Dietetics and Nutrition by Katelynn Burghardt, RDN, with support from Debra Sullivan, Ph.D., RD, Department Chair and Midwest Dairy Council Professor in Clinical Nutrition.

This cookbook was made possible with the generous support of Otsuka America Pharmaceutical, Inc.

Published by the PKD Foundation



**PKD FOUNDATION**  
Polycystic Kidney Disease

1.800.PKD.CURE  
[pkdcure.org](http://pkdcure.org)

Produced by Ascend Integrated Media  
6710 W. 121st St., Suite 100  
Overland Park, KS 66209  
[ascendmedia.com](http://ascendmedia.com)

Recipe photography by Wheat Photography  
115 W. 18th St., Suite 200  
Kansas City, MO 64108  
[wheatphoto.com](http://wheatphoto.com)

# Cooking **well**

Delicious,  
everyday recipes  
for the PKD family

## **ABOUT THE PKD FOUNDATION**

The PKD Foundation is dedicated to finding treatments and a cure for polycystic kidney disease (PKD) to improve the lives of those it affects. We do this by funding research and promoting education, advocacy, support and awareness.

Our vision is that one day no one will suffer the full effects of PKD. Visit [pkdcure.org](http://pkdcure.org) to learn more.



**PKD FOUNDATION**  
Polycystic Kidney Disease

## A Note From the Author



After becoming a dietitian in 2011, I moved to Kansas City, Missouri, to complete my Ph.D. in Medical Nutrition Science at the University of Kansas Medical Center. As a first-year Ph.D. student, I was given the task of selecting a research topic for my dissertation, a choice that may ultimately define my career. At that time, I was working as a pediatric nephrology dietitian at Children's Mercy Hospital in Kansas City, so doing research in kidney disease was of significant interest. Fortunately, I was approached by physician-scientist and PKD Foundation co-founder Jared Grantham, M.D., to design a diet for patients with PKD. Needless to say, with my background in clinical nutrition, I was intrigued by the possibility of designing a diet based on scientific evidence that would benefit patients.

Dr. Grantham thought that because certain dietary elements may potentially affect the kidneys of individuals with PKD, a special diet should be designed and prescribed early in the course of the disease. Together, we developed a diet that would limit potentially harmful dietary elements, like excessive intakes of sodium and protein, and add more elements that may protect the kidney, like fruits, vegetables and fluids. This cookbook is not just a cookbook of delicious, healthy foods, but is based on the research of many scientists working to understand how diet may be used as a therapy for PKD patients. These recipes adhere to a set of dietary goals based on that research.

After finishing my doctorate in 2015, I relocated to Groningen, the Netherlands, to complete my post-doctorate training at the University Medical Center Groningen in its Expertise Center for PKD research. My work continues to be centered on understanding how diet may affect cyst growth and kidney damage in PKD and how making dietary changes may help protect the kidney.

It is my hope that this cookbook provides those with PKD, who are looking to play an active role in their disease, a helpful hand in understanding how to change their diet, with recipes to suit.

kindest regards (Met vriendelijke groet),  
Jacob Taylor, Ph.D., RD, LD

# Table of Contents

## NUTRITIONAL GUIDELINES

For Individuals Living With ADPKD	6
For Individuals on Dialysis	16
For Individuals With Kidney Transplants	22
For Children at Risk of Having ADPKD and Individuals With ARPKD	26

## RECIPES

Appetizers	32
Soups and Salads	46
Sides	62
Entrees	74
Desserts	88
Spices, Spreads and Rubs	102

## REFERENCE

Glossary of Terms and Topics	109
References	114
Index	117



**PKD FOUNDATION**  
Polycystic Kidney Disease

1.800.PKD.CURE  
pkdcure.org



# Nutrition Guidelines for **Individuals** Living With **ADPKD**

Your goal as someone living with PKD should be to maintain a healthy lifestyle and to protect your kidneys as best you can. Autosomal dominant PKD (ADPKD) researchers are currently examining the potential role that diet may play in protecting your kidneys. Several dietary factors have been shown to be harmful to individuals with PKD.<sup>1-11</sup> For example, eating too much sodium and animal protein may be harmful to your kidneys, along with having low HDL cholesterol levels (HDL is the good cholesterol) and carrying around extra body weight.<sup>1,2,8,11</sup> Caffeine should also be limited.<sup>9,10</sup> On the other hand, drinking water and eating fruits and vegetables may play a role in protecting your kidney.<sup>2-7,12-14,15</sup>

The following is a general set of guidelines to help you eat healthy while living with PKD. Doing these things will help cut down the workload of your kidneys and help preserve your kidney function. Your doctor and/or dietitian may prescribe different guidelines depending on your health and lab values. See page 110 for more on the difference between a dietitian and nutritionist.

## **GOALS OF LIVING WITH A PKD DIET**

- Suppress a hormone known as vasopressin, which causes cysts to grow.<sup>6</sup>
- Keep your cholesterol levels in normal range, especially HDL (good cholesterol).<sup>1</sup>
- Maintain a healthy body weight. <sup>1</sup>
- Reduce the dietary acid load of your diet.<sup>2,3,12-15</sup>
- Limit caffeine consumption.<sup>9,10</sup>
- Prevent diabetes and prevent or control high blood pressure. Both of these conditions are harmful to the kidneys.<sup>16</sup>





## GENERAL DIET GUIDELINES

### MAINTAIN A HEALTHY WEIGHT

One goal to strive for when trying to be healthy is to maintain a healthy body weight. For an individual without PKD, a healthy body weight is measured by determining your body mass index (BMI). Your BMI is simply a measure of how much you weigh compared to how tall you are. Having a BMI between 18.5 and 24.9 is ideal. However, because many PKD patients have large, heavy kidneys due to cysts, it is not always appropriate to use body weight or BMI as a measurement tool. PKD patients often refer to this as having a “PKD belly,” in reference to having a large midsection caused by their cystic kidneys. There is little you can do to make this go away, but adopting a healthy diet and daily exercise will ensure you are as healthy as you can be. Rather than focusing solely on the number on the scale, we encourage you to stay as healthy as possible by focusing on healthy eating behaviors and exercising regularly. If you would like to have a target weight or BMI, please work with your physician or dietitian to come to an agreed-upon number. Read on to learn nutrition guidelines created for PKD patients to address your unique needs.

### SODIUM

High-sodium diets raise levels of vasopressin, a hormone that may be harmful to cystic kidneys, in the body.<sup>16</sup> People primarily consume sodium in the form of salt. The majority of sodium — approximately 75 percent — in the American diet comes from processed and packaged foods.<sup>18</sup> When at the grocery store, remember to check the nutrition label for sodium content. If you’re eating out, most restaurants will have their nutrition information posted online. This information will list sodium content for all of their food items. **You should limit yourself to**

<b>Nutritional Facts</b>	
Serving Size 1 cup	
Servings Per Container 2	
<b>Amount Per Serving</b>	
<b>Calories</b> 250	Calories from Fat 110
<b>% Daily Value*</b>	
<b>Total Fat</b> 12g	<b>18%</b>
Saturated Fat 3g	<b>15%</b>
Trans Fat 3g	
Cholesterol 30mg	<b>10%</b>
<b>Sodium</b> 360mg	<b>15%</b>
<b>Total Carbohydrate</b> 31g	<b>10%</b>
Dietary Fiber 0g	<b>0%</b>
Sugars 5g	
<b>Protein</b> 5g	
Vitamin A	4%
Vitamin C	2%
Calcium	20%
Iron	4%
*Percent Daily Values are based on a 2,000 calorie diet. Your Daily Values may be higher or lower depending on your calorie needs.	
	Calories: 2,000 2,500
Total Fat	Less than 65g 80g
Saturated Fat	Less than 20g 25g
Cholesterol	Less than 300mg 300mg
Sodium	Less than 2,400mg 2,400mg
Total Carbohydrate	300g 375g
Dietary Fiber	25g 30g



**2,000 milligrams of sodium per day.** A general guideline to keep from going over this would be to limit meals to 500 milligrams of sodium and snacks to 200 milligrams. This will not only help keep vasopressin levels low, it also may help keep your blood pressure in the normal range, making it a dietary component worth paying attention to.

### PROTEIN

**Overall, protein should be limited based on your body weight.**<sup>1,2,8,11</sup>

To calculate your protein needs, all you need to know is your height in inches, which can be put into the following formula:

#### **Daily Protein Needs (grams per day)**

$$(21.7 \times (\text{height in inches})^2 / 703) \times 0.36$$

*Example:* If Bob is 5 feet, 10 inches (70 inches), his protein needs would be calculated as follows:

**Step 1:**  $(70 \text{ inches})^2 = 4,900$

**Step 2:**  $(21.7 \times 4900) = 106,330$

**Step 3:**  $(106,330 / 703) = 151$

**Step 4:**  $151 \times 0.36 = 54$  grams per day

Bob should try to consume 54 grams of protein each day.

Following this calculation will ensure that you meet your protein needs while preventing excess protein intake from harming your kidneys. Animal-source proteins (chicken, pork, beef, fish, eggs, cheese/milk, etc.) can certainly be part of your diet, but **should be limited to approximately three-ounce portions at meal times or no more than six ounces per day** (weight is after cooking). Try to include fatty fish, like salmon, as your protein choice at least two times per week.

### POTASSIUM

Potassium may play an important role in protecting the kidneys by reducing your dietary acid load.<sup>2,3,12-15</sup> Not to be confused with acidic foods (like lemons), dietary acid load is a measure of the





amount of protein-to-potassium an individual eats.<sup>19</sup> Basically, dietary acid load is high in individuals whose diets consist of moderate to large amounts of meat (protein) with minimal intake of fruits and vegetables (potassium), making fruits and vegetables a critical component of your diet. See page 13 for more information about dietary acid load. Potassium is also an important nutrient for keeping your blood pressure in the normal range.<sup>20</sup> **The recommended potassium intake is 4,700 milligrams a day.**<sup>21</sup> As your kidney function declines, you may need to start restricting potassium; however, your physician will notify you of this at that time. To meet your daily requirements, you should aim for eating at least one fruit or vegetable at every meal, and make sure at least some of those are green and leafy, such as spinach, kale or broccoli.

## FLUIDS

Drinking fluids is very important for keeping your kidneys healthy. Like reducing your sodium intake, staying hydrated will keep your vasopressin levels low.<sup>5-7</sup> For adults, **at least three liters (100 ounces) a day is recommended and should be spread out over the course of the day.**<sup>5</sup> These fluids should **primarily be from non-caffeinated beverages.**<sup>9</sup> See page 11 for additional information about water and ADPKD.



Serving Equivalent	
Dairy or Substitute Products	Amount That Counts as a Cup in the Dairy Group
Milk (choose fat-free or low-fat milk)	1 cup milk
	1 half-pint container milk
	½ cup evaporated milk
Yogurt (choose fat-free or low-fat yogurt)	1 regular container (8 fluid ounces)
	1 cup yogurt
Cheese (choose reduced-fat or low-fat cheeses)	1 ½ ounces hard cheese (cheddar, mozzarella, Swiss, Parmesan)
	⅓ cup shredded cheese
	2 ounces processed cheese (American)
	½ cup ricotta cheese
	2 cups cottage cheese
Milk-based desserts (choose fat-free or low-fat types)	1 cup pudding made with milk
	1 cup frozen yogurt
	1 ½ cups ice cream
Almond, rice or soy milk fortified with calcium	1 cup calcium-fortified milk alternative
	1 half-pint container calcium-fortified milk alternative

SOURCE: CHOOSEMYPLATE.GOV/DAIRY

## CALCIUM

Most adults need to consume 1,000 milligrams of calcium daily.<sup>21</sup> This equates to three servings of dairy daily. For women over 51 years old and men over 70 years old, four servings of dairy daily are recommended.<sup>21</sup> A serving of dairy is one ounce of cheese or eight ounces of

milk or yogurt. Many dairy alternatives, such as almond milk, are also fortified with calcium.

## CAFFEINE

Caffeine has been shown to have the potential to accelerate cyst growth in ADPKD.<sup>9</sup> No recommendations have been made on the exact amount of caffeine individuals with ADPKD should limit themselves to each day. At the present time, while **caffeine should be limited in individuals living with ADPKD**, the consumption of a single cup of coffee (eight ounces equals 70 to 200 milligrams of caffeine), two cups of tea (16 ounces equals 30 to 140 milligrams of caffeine), or one diet soda (20 ounces equals 58 to 100 milligrams of caffeine) daily may be acceptable.<sup>10</sup> Decaf or caffeine-free versions of drinks are more appropriate, and if possible should be selected in place of a caffeinated beverage. **Caffeine-containing supplements, energy shots and energy drinks should be avoided due to their high caffeine content.**



## CALORIES

**Calorie intake should be appropriate to maintain a healthy body weight.** For men who are sedentary, the recommended daily calorie intake is between 2,000 and 2,600 calories, and for women who are sedentary, it is between 1,600 and 2,000.<sup>22</sup> These are general guidelines and may not be suitable for every individual. Watching how your calorie intake affects your body weight is the best way to determine whether you are eating too few or too many calories based on your goals.

## EXERCISE

As with any healthy diet, exercise should be a part of a broader healthy lifestyle. **Thirty minutes of exercise per day at least five days per week** will help keep your weight under control and may help improve HDL cholesterol (good cholesterol).<sup>23</sup> If you smoke, quitting may help increase your HDL cholesterol, as well as help improve your heart and lung health.<sup>24</sup>



## WATER AND ADPKD

Drinking plenty of water has many potential benefits, including reducing the risk of kidney stones and preventing dehydration, which can worsen kidney function. To properly regulate fluid balance in our bodies, we depend on a hormone called vasopressin to conserve water when we are dehydrated and excrete water when we have too much. Unfortunately, high vasopressin levels (which occur when we don't get enough fluid) have been linked to many diseases, including high blood pressure, heart and kidney disease, and a particularly harmful effect in ADPKD. It is known that drinking more fluid can reduce vasopressin and slow the progression of ADPKD. However, drinking enough fluid each day can be difficult for patients to maintain in real life. Part of the difficulty in keeping vasopressin levels low throughout the day by drinking water is that people eat too



much salt and protein, raising vasopressin levels to help conserve water and maintain normal fluid balance. For the most part, daily intake of water, sodium and protein is not determined by physiologic requirements, but is more a function of dietary preferences and cultural influences. Healthy adults have an average daily fluid intake of one to three liters (34 to 101 ounces), with considerable individual variation. Thirty-three percent of this comes from a combination of water in food and your body producing water when it burns fat for energy, while the other 66 percent comes from the fluid you drink. Similarly, of the high solute ingested in the typical American diet, 75 percent of the solutes come from sodium, and the remaining 25 percent of solutes come from protein intake.<sup>25</sup> Solute is a dietary compound that stimulates vasopressin secretion.

To address the challenges associated with a high fluid intake, a novel dietary intervention that lowers the amount of water needed to suppress vasopressin was developed. In a randomized controlled trial conducted at Tufts Medical Center on 34 patients with ADPKD, a diet-based intervention lowered vasopressin and reduced the amount of water patients needed to drink to achieve a reduction in vasopressin.<sup>26</sup> This intervention is based on a two-step approach of combining a diet low in protein and salt (low osmolar diet) with individualized adjusted water intake. Water intake was based on how much salt and protein were consumed. In this trial, individuals ingested 2.6 liters (88 ounces) of water per day on average, which significantly reduced vasopressin in patients with ADPKD. The average American consumes 1.2 liters per day.<sup>27</sup> It is important to note that the amount of water needed in patients with ADPKD is directly related to the amount of salt and protein eaten each day. Thus, the more salt and protein you eat, the more water you need to drink to achieve the maximum benefit. Water in this context refers to tap and bottled water only, with no additions. Individuals are encouraged to drink additional fluid, if desired. This includes, but is not limited to, vitamin water, tea and juice. Ideally, water intake should be distributed throughout the waking hours.





In summary, drinking plenty of water offers a safe, easily tolerated and affordable approach that can be adopted in the early stages of ADPKD to help keep your kidneys healthy.

*Dr. Osama Amro would like acknowledge Dr. Ronald Perrone for his role as a supervisor of the dietary and water study at Tufts Medical Center.*

### **DIET AND PKD: A PILOT INTERVENTION STUDY**

We have discussed the benefits of drinking fluid and eating less sodium and protein. Although these three dietary factors are important for individuals living with ADPKD, the additional dietary factor to be discussed is dietary acid load (see pages 8-9 for dietary acid load intro and guidelines). As you have already learned, dietary acid load, put simply, is the relationship between the amount of animal protein you consume and the amount of fruits and vegetables you eat. Animal proteins are metabolized into acids when eaten, which appear to be harmful to kidney function. Fruits and vegetables are metabolized into bases and may protect the kidneys.<sup>2,3,12-15</sup> These findings led researchers to examine whether individuals living with ADPKD may benefit from following a low-sodium, lower-protein diet supplemented with additional fruits, vegetables and fluids. Additionally, they wanted to determine whether this type of diet was feasible for individuals.

A pilot study was conducted in individuals with ADPKD at the University of Kansas Medical Center in Kansas City, Kansas.



Twelve adults with ADPKD were enrolled in a five-week study to determine whether individuals could follow a diet lower in sodium and protein and higher in fruits, vegetables and water.<sup>28</sup> Individuals consumed their usual diet for one week and the Kansas PKD diet for four weeks. Eleven individuals completed the study, and results showed a 36 percent reduction in sodium intake, a 28 percent reduction in protein intake, a 99 percent decrease in dietary acid load and a 42 percent increase in fluid intake. These findings were consistent with what was collected in urine samples during the study. Participants were not only successful following the Kansas PKD diet, but most participants felt “somewhat confident” or “very confident” in their ability to follow the Kansas PKD diet. This is not to say that individuals found no difficulty in following the diet, but that these difficulties were things they felt confident in being able to manage. Individuals reported that reaching the goal

fruit and vegetable intake each day was difficult, yet participants increased their fruit and vegetable intake three-fold over what they had been eating. Eating out was reported as the most difficult time to adhere to the diet; however, you can still enjoy these things if you know what nutrition information to look for and how best to order at a restaurant. Look for lower-sodium meals (less than 500 milligrams) that don't contain too much protein (less than 25 grams), and don't skip the fruits or vegetables. Most restaurants' nutrition information can be provided by the restaurant, found online or accessed on your smartphone.

The findings from this study indicate that individuals living with ADPKD can follow a relatively complex diet prescription that may help protect their kidneys, at least over the course of one month. Plus, individuals were able to do so with only modest stressors. The results of this study indicate that for those individuals looking to play an active role in their disease, the dietary intervention





used here (and detailed in this cookbook with recipes that fit these dietary guidelines) can be followed by individuals safely, confidently and started as early as childhood. Family members without PKD also can follow this diet, as it adheres to healthy eating guidelines.

### **VEGETARIAN OR VEGAN DIETS AND ADPKD**

Many individuals living with PKD may ask: If certain dietary behaviors are bad, shouldn't I just avoid them altogether? One topic that crops up frequently is about switching from a diet that includes meat to a vegetarian or vegan diet. To clarify the difference, vegetarians exclude most meats from the diet, but may still consume dairy, eggs or fish (or any combination of the three), while vegans exclude animal products of any kind from their diet (no meat, dairy, eggs, fish or any products or foods that include any animal-based products, including foods with gelatin). When followed correctly, these types of diets have many health benefits, such as reducing risk of cancer, cardiovascular disease, obesity, high cholesterol levels, high blood pressure, type 2 diabetes and even improving quality of life.<sup>29-33</sup> In addition to these benefits, it may even be protective of kidneys in individuals with PKD. Currently, no studies have been conducted to determine whether a vegetarian or vegan diet is more beneficial than a diet low in animal protein. At this time, either limiting meat intake or following a vegetarian or vegan diet can be recommended for individuals living with PKD. However, it is important to understand that animal protein foods contain essential nutrients that, when eliminated from the diet, can put you at risk for developing nutrient deficiencies that lead to health problems (including iron, zinc and B12 deficiency, to name a few). **If you are considering switching to a vegetarian or vegan diet, consult your physician and dietitian first, so that he or she can properly instruct you on how to make the appropriate dietary changes to meet your nutrient needs or whether certain supplements may be necessary.** Your physician may want to monitor you more closely initially to make sure that everything is being done safely and appropriately.



# Nutrition Guidelines for **Individuals** on **Dialysis**

**W**ithout functioning kidneys, certain diet-associated waste products may build up in your blood. Watching what you eat and drink will help prevent this from occurring. Your doctor and dietitian should notify you about what you need to do to change your diet. Your goal should be to stay healthy by making sure your labs are in the normal range. **While continuing to restrict sodium is appropriate, other dietary recommendations mentioned in the “Nutrition Guidelines for Individuals Living With ADPKD” section**



Look for this icon on recipes for **dialysis-friendly** options.

**MAY NOT be appropriate.** Many patients require a potassium restriction (due to high potassium levels in the blood) or must restrict fluid intake (due to producing less urine) on dialysis. You also need more protein in your diet since small amounts of protein may be filtered out during dialysis. Your

doctor and dietitian will instruct you to do this by eating a little more meat or by prescribing you a nutritional supplement. For vegetarians or vegans, they will provide other alternatives. It is important for you to understand how to make these changes by making simple alterations to meals, and why doing these things is important to achieving optimal health.

The following is a general set of guidelines to help you eat healthy while on dialysis. Your doctor and dietitian may prescribe different guidelines depending on your health and lab values. See page 110 for more on the difference between a dietitian and nutritionist.

## **GOALS OF THE DIALYSIS DIET**

- Keep serum potassium in normal range (3.5 to 5.5 millimoles per liter).<sup>34</sup>
- Keep serum phosphorus levels in normal range (3.5 to 5.5 milligrams per deciliter).<sup>35</sup>





- Keep your fluid weight gain between dialysis treatments at an appropriate level (usually less than two kilograms or 4.4 pounds).<sup>36</sup>
- Maintain an appropriate nPCR or nPNA level (minimum of 0.8; optimal is 1 to 1.4).<sup>37</sup>
- Keep your blood pressure in normal range, so your heart will be healthy (140/90 mmHg and 130/80 mmHg, pre- and post-dialysis, respectively).<sup>36</sup>

Doing these things will help keep you as healthy as possible.

## GENERAL DIET GUIDELINES

### SODIUM

Watching your sodium intake while on dialysis will be important for maintaining a healthy blood pressure. Sodium will not only cause you to be thirsty and drink more fluid, but your body will also retain more fluid and raise your blood pressure. Sodium is primarily consumed in the form of salt. The majority of sodium in the American diet comes from processed and packaged foods.<sup>18</sup> When at the grocery store, remember to check the nutrition label for sodium. **You should limit yourself to 2,000 milligrams of sodium per day.** A general guideline to keep from going over this would be to limit meals to

<b>Nutritional Facts</b>			
Serving Size 1 cup			
Servings Per Container 2			
<b>Amount Per Serving</b>			
<b>Calories 250</b> Calories from Fat 110			
			% Daily Value*
<b>Total Fat</b>	12g		<b>18%</b>
<b>Saturated Fat</b>	3g		<b>15%</b>
<b>Trans Fat</b>	3g		
<b>Cholesterol</b>	30mg		<b>10%</b>
<b>Sodium</b>	360mg		<b>15%</b>
<b>Total Carbohydrate</b>	31g		<b>10%</b>
<b>Dietary Fiber</b>	0g		<b>0%</b>
<b>Sugars</b>	5g		
<b>Protein 5g</b>			
<b>Vitamin A</b>			<b>4%</b>
<b>Vitamin C</b>			<b>2%</b>
<b>Calcium</b>			<b>20%</b>
<b>Iron</b>			<b>4%</b>
*Percent Daily Values are based on a 2,000 calorie diet. Your Daily Values may be higher or lower depending on your calorie needs.			
		Calories:	2,000    2,500
<b>Total Fat</b>	Less than	65g	80g
<b>Saturated Fat</b>	Less than	20g	25g
<b>Cholesterol</b>	Less than	300mg	300mg
<b>Sodium</b>	Less than	2,400mg	2,400mg
<b>Total Carbohydrate</b>		300g	375g
<b>Dietary Fiber</b>		25g	30g



500 milligrams of sodium and snacks to 200 milligrams. Limiting packaged and processed foods in the diet will also help you limit the additional phosphorus they often have. It is also important to note that reduced sodium foods, particularly soups, may add potassium chloride (NuSalt) in its place. If you are on a potassium restriction, it is important to pay attention to this, as it should be avoided.

### PROTEIN

The National Kidney Foundation's Kidney Disease Outcomes Quality Initiative (KDOQI) guidelines recommend 1.2 to 1.3 grams per kilogram body weight per day (0.55 to 0.59 grams per pound of body weight per day).<sup>37</sup> This is more than is recommended for patients living with PKD or after having a kidney transplant due to the fact that dialysis removes some protein from the body. It usually is recommended that half of your protein intake comes from animal-source proteins (chicken, pork, beef, fish, eggs, cheese/milk, etc.).

However, this is still only about six ounces of meat per day. If you are on hemodialysis, your doctor or dietitian may alert you to your nPCR level, which measures whether or not you have enough protein in your diet. Alternatively, if you are on peritoneal dialysis, your doctor or dietitian may alert you to your nPNA level. For individuals who are vegetarian or vegan, your dietitian will discuss with you alternative ways to make sure you meet your protein needs.



## POTASSIUM

Potassium is of great importance before dialysis and after a kidney transplant. **While on dialysis, you may need to limit your intake of potassium.** This doesn't mean you have to stop eating fruits and vegetables, but you may need to be more selective. Apples, pears and peaches may be fine, but oranges and bananas should be eaten sparingly. For vegetables, green beans, cauliflower and onions are all low in potassium, while potatoes and tomatoes should be eaten less often. Some high-potassium foods, such as potatoes, may be fine if you cook them in a particular way. To reduce the potassium content of potatoes, dice or slice into small cubes or slivers. Then, boil them for 10 minutes. Cutting them smaller will remove more potassium.

Doing this can result in removing 50 percent of the potassium in potatoes cut into small cubes and 75 percent for potatoes sliced into thin slivers.<sup>38</sup> For most patients, potassium should be kept to 2,000 to 3,000 milligrams per day while on dialysis. However, patients on peritoneal dialysis may not need to be as restrictive, since they receive dialysis every day.



### **HOW TO REDUCE THE POTASSIUM CONTENT OF POTATOES, SWEET POTATOES, CARROTS, BEETS, WINTER SQUASH AND RUTABAGAS**

**Step 1:** Slice vegetables into cubes no more than ½ inch thick or cut into thin slivers. If peeling prior to slicing, place the vegetables in cold water to prevent darkening.

**Step 2:** Bring water to a boil and cook for 10 minutes.

**Step 3:** Pour off water (discard) and enjoy.



## FLUIDS

While on dialysis, you often have a decrease in the amount of urine you make each day, or you may not make any urine at all. This means that you may need to limit the amount of fluid in your diet to prevent storing a large amount of fluid on your body that must then be removed by dialysis. Your doctor or dietitian may note your body weight between dialysis treatments to measure how much fluid you are retaining. If you are retaining too much fluid, **you may need to limit your fluid intake to less than two liters per day (67 ounces)**. However, your doctor or dietitian will notify you of your specific goal. Fluid-based foods such as soups and smoothies also count toward your daily total.

## PHOSPHORUS

If you eat too much phosphorus while on dialysis, it can harm your heart and bones. Phosphorus can be found in many foods, including meats, dairy, whole grains and dark colas. Phosphorus



is a preservative and is found in most packaged and processed foods. Fortunately, it is possible to reduce the phosphorus content of the meat you consume. Similar to reducing potassium in vegetables, meat should be cut into thin slices and boiled for at least 10 minutes (30 minutes is preferred). Doing this can result in the removal of 50 percent of the phosphorus by boiling for 10 minutes and 65 percent of the phosphorus by boiling for 30 minutes.<sup>39</sup> For most patients, phosphorus should be limited to no

more than 800 to 1,000 milligrams per day. **You should aim to keep your serum phosphorus levels less than 5.5 milligrams per deciliter.**<sup>35</sup> Your physician also may prescribe you phosphorus binders to be taken with meals to help prevent phosphorus absorption.

## HOW TO REDUCE THE PHOSPHORUS CONTENT OF MEAT

**Step 1:** Slice meat into thin strips.

**Step 2:** Bring water to a boil. Boil for at least 10 minutes (30 minutes is preferred).

**Step 3:** Pour off water (discard) and enjoy.

## CALCIUM

While on dialysis, you should aim for 1,000 milligrams of calcium per day.<sup>21</sup> Since you may have to limit your intake of milk and cheese because of the phosphorus content, your physician may prescribe a calcium supplement to be taken with or away from meals to help meet your needs. Keep in mind that milk substitutes such as soy milk may contain phosphorus additives, so make sure you check the label before using them as a source of calcium.

## CALORIES

Eating enough calories to get proper nourishment while maintaining a healthy body weight while on dialysis is important. The KDOQI guidelines recommend getting 35 calories per kilogram body weight (16 calories per pound) for people under 60 years of age and 30 to 35 calories per kilogram body weight (13.6 to 16 calories per pound) for people over 60 years of age.<sup>37</sup>



# Nutrition Guidelines for **Individuals With Kidney Transplants**

**F**or individuals who have received a transplant, your diet will resemble the foods you ate prior to dialysis, but for different reasons. Watching your sodium intake will be important for keeping your blood pressure under control, while drinking sufficient amounts of water will be important for making sure your kidney gets plenty of blood flow. Fruits and vegetables will still be important for reducing your dietary acid load. Limiting meat will also be important.<sup>40</sup> The dietary restrictions you had if you were on dialysis will ease up once your kidney function improves, so it is important to make the appropriate changes to your diet. Your doctor and dietitian will help you transition from the diet you may have followed on dialysis to your new diet. See page 110 for more on the difference between a dietitian and nutritionist.

## **GUIDELINES FOR AFTER YOUR KIDNEY TRANSPLANT**

### **PREVENT EXCESS WEIGHT GAIN**

After having a kidney transplant, patients often feel better, and without the strict dietary restrictions that are often necessary if on dialysis, people tend to overeat and gain weight.<sup>41</sup> Some medications prescribed to individuals with kidney transplants also can cause weight gain.<sup>42</sup> Gaining weight can increase your risk for several chronic diseases, including hypertension and diabetes. This can be bad for your new kidney. It is important to remember to avoid overeating, watch your salt and sugar intake and exercise regularly if your doctor has cleared you to do so.





## KEEP YOUR BONES HEALTHY

Kidney disease can cause your bones to weaken. Calcium is responsible for keeping bones strong and an important part of everyone's diet. **Ideally, calcium can be obtained by consuming three servings of dairy per day.** However, other foods may be

fortified with calcium, including orange juice and milk substitutes. These foods may be consumed as an alternative to dairy products to meet your calcium needs. Check labels to see how much calcium is contained per serving. If getting calcium from food isn't possible, talk to your doctor and dietitian about taking a calcium supplement to help meet your needs.

Serving Equivalent	
Dairy or Substitute Products	Amount That Counts as a Cup in the Dairy Group
Milk (choose fat-free or low-fat milk)	1 cup milk
	1 half-pint container milk
	½ cup evaporated milk
Yogurt (choose fat-free or low-fat yogurt)	1 regular container (8 fluid ounces)
	1 cup yogurt
Cheese (choose reduced-fat or low-fat cheeses)	1 ½ ounces hard cheese (cheddar, mozzarella, Swiss, Parmesan)
	⅓ cup shredded cheese
	2 ounces processed cheese (American)
	½ cup ricotta cheese
	2 cups cottage cheese
Milk-based desserts (choose fat-free or low-fat types)	1 cup pudding made with milk
	1 cup frozen yogurt
	1 ½ cups ice cream
Almond, rice or soy milk fortified with calcium	1 cup calcium-fortified milk alternative
	1 half-pint container calcium-fortified milk alternative

SOURCE: CHOOSEMYPLATE.GOV/DAIRY

## HERBAL SUPPLEMENTS

After having a kidney transplant, you likely will be on medications that suppress the immune system. Suppressing the immune system is important so that the body doesn't attack the new kidney. For this reason, you should **avoid taking herbal supplements to help boost the immune system.** These supplements interfere with your other post-transplant medications. Supplements that may be beneficial after having a kidney transplant include fish oil, approved multivitamins and calcium. Check with your doctor and dietitian if you are considering one of these supplements.





## FOOD SAFETY

Food safety is important for everyone, but for individuals with kidney transplants who are often on immune-suppressing medications, it is of the utmost priority. **Following food safety protocols after having a kidney transplant is very important.** [Foodsafety.gov](https://www.foodsafety.gov) uses the simple mantra of “clean, separate, cook, chill” to help you understand the four necessary steps to keeping food safe.

### CLEAN

Washing hands is a key component to keeping food safe. Washing surfaces, utensils and cutting boards is necessary to prevent the spread of bacteria, especially when they have come into contact with raw meat. Finally, it is important to wash fruits and vegetables before eating. This includes fruits and vegetables with rinds, such as watermelon, lemons and cantaloupe. Although you may not eat the rind, the knife will spread the

bacteria from the outer surface into the core when cutting. Like us, bacteria eat the fruit for energy, which results in them rapidly increasing in number and potentially making you sick.

### SEPARATE

Keep uncooked meat away from other foods. While at the grocery store, keep the meat in a separate part of your grocery cart. Just because the meat is wrapped in packaging doesn't mean it won't leak onto your fresh fruits and vegetables. When preparing raw meat you should use a separate cutting board with separate utensils. As always, don't forget to wash your hands. Finally, meat should always be packaged neatly and stored at the bottom of the refrigerator to prevent leakage onto other foods.




## COOK

Cook all meats to the appropriate temperatures. To do this, make sure you have a food thermometer available. The food thermometer should be placed in the thickest part of the meat, and the temperature checked to make sure the entire cut of meat is cooked thoroughly. Once cooked, keep foods above 140°F to ensure they don't go into the danger zone where bacteria can rapidly multiply. When eating leftovers, microwave foods to 165°F and check with a food thermometer. Make sure the food is stirred while microwaving to ensure that the food is heated evenly.

## CHILL

After eating, make sure the food is stored properly. Food should be wrapped up and chilled within two hours after cooking is complete. See the food chart at [fda.gov](http://fda.gov) to determine how many days foods can safely be left in the refrigerator and freezer.

Especially during summer months, you should be careful about food at barbecues and picnics. When not stored at the proper temperatures, foods that should be chilled, such as potato and macaroni salads, and foods that need to remain at or above 140°F, such as meat, may be sitting out without the proper chilling or heating devices, making them a high risk for bacterial growth. If you are ever uncertain, it is best to avoid these types of foods.



# Nutrition Guidelines for **Children at Risk of Having ADPKD** and **Individuals With ARPKD**

## **ARPKD OR ADPKD**

Individuals living with PKD may have one of two forms: autosomal recessive PKD (ARPKD) or autosomal dominant PKD (ADPKD). ARPKD is faster-progressing, and the onset of symptoms is generally in childhood. ADPKD may not present symptoms until adults are middle-aged. Most of the diet research has been conducted in ADPKD patients or animal models that more closely resemble ADPKD. However, the dietary factors studied in these research trials ultimately target certain hormones or biochemicals known to cause cyst growth. Although clinical trials have not been performed to examine whether these same dietary factors influence disease progression in ARPKD, individuals living with ARPKD may benefit from making these same dietary changes.

## **FEEDING A CHILD WHO HAS PKD**

Many parents may not know whether their child has PKD or not. Parents may ask themselves whether their child should make diet changes, just in case, or whether the diet may be harmful or affect their child negatively if he or she ends up not having PKD. Fortunately, the dietary factors discussed in the *Nutrition Guidelines for Individuals Living With ADPKD* section of this cookbook are all healthy dietary behaviors that can be followed by anyone in the family, regardless of whether or not he or she has PKD. Therefore, special foods do not need to be prepared for a single individual, but can be prepared for the entire family. It is safe for your child to follow a PKD-friendly diet, even if he or she ends up not having PKD.



Ultimately, eating healthy as a child will promote lifelong healthy dietary behaviors and may lower his or her risk for certain chronic diseases, such as high blood pressure or diabetes, during adulthood.

It is also important to know that making PKD-friendly dietary choices should begin as soon as children begin eating solid foods, rather than waiting until they have symptoms decades later. This is because PKD (particularly ADPKD) is a very slow-progressing disease that takes years until the onset of symptoms. If your child has ADPKD, they have it from birth. Yet, symptoms aren't noticeable until decades later when the disease has already caused a lot of damage to the kidney. If symptoms do appear during childhood, consult your family doctor or family nephrologist to determine what should be done next for your child. Your physician or dietitian can instruct you on whether changes to your child's diet need to be made that differ from the dietary guidelines in this cookbook. See page 110 for more on the difference between a dietitian and nutritionist.

#### AGES 1 THROUGH 4

Exposure to new foods is key at this age. **Children may not be open to trying foods or liking certain foods until they have been exposed to it up to 15 times.**<sup>45,46</sup>

Parents may only offer a food to a child once and, if their child doesn't want to try it or appears to dislike it, the food is never offered again, leaving the parent to believe their child is a picky eater. In addition to trying a variety of foods, make sure to offer fruits and vegetables at every meal. This will help ensure that your child gets enough potassium. Your child should consume fluid, primarily water, over the course of the day. Have a sippy cup around for the child to drink out of once they are old enough. While protein is important for growth and development, meat should be limited to meal times to avoid eating excess protein. Finally, buying fewer processed foods and cooking at home more frequently will reduce your child's sodium intake and help keep vasopressin levels low, which should help protect their kidneys.





### AGES 5 THROUGH 12

Once children reach school age, their eating behaviors are more influenced by peers. To continue to promote healthy eating, sending lunch from home may ensure they get the food they need to keep their kidneys healthy. Some children at this age may be more interested in eating a school-provided lunch. Parents can look at the meals prior to the school day and decide which days are best for the child to eat in the cafeteria and which days food should be sent from home. You or your physician may even request a special diet from the school that will help make sure your child gets the right foods.

In addition to eating healthy, children at this age may start being involved in sports. It is important that your child's doctor has approved exercise and that there are no potential issues with your child playing the sport of his or her choice. Some contact sports such as football may not be recommended. When engaging in sports, it is important to make sure children stay hydrated. In most circumstances, water is the preferred source of fluids for your child during exercise. Sports drinks are often loaded with sodium. Unless your child is an elite athlete training for hours a day or playing in extreme temperatures, water should be sufficient for hydration. Children also may benefit from carrying a water bottle to classes



throughout the day. Doing this will help ensure they stay hydrated and that vasopressin levels are suppressed over the course of the day.

Given your child's busy schedule during these years, preparing meals every day may not be possible. Often, fast food is of great convenience, even if it doesn't have the healthiest options. Fast food establishments should have nutrition facts available in the restaurant or online (accessible via smartphone). It would be a good idea to look up a few "best" options at the most common restaurants your child likes. One thing you can do to make fast food healthier is to request fresh, unsalted fries. This will greatly reduce the amount of sodium in your child's meal. You should pay particular attention to the sodium and protein contents of the meal. **Meals should primarily contain less than 500 milligrams of sodium and no more than 10 grams of protein for children between 5 and 8 years of age, no more than 15 grams of protein for children 9 and 12 years of age and no more than 20 grams of protein for children 13 and 18 years of age.**

Finally, involving children in the preparation of dinner will be important to increase the chances that they will try new foods at meal times. Involvement can be limited to doing simple tasks, such as setting the table or getting small items from the refrigerator. As children get older and have mastered the safe use of utensils, they may be able to help with more difficult tasks related to food preparation. **This will increase the chances of them trying new foods and teach them cooking skills that will be useful throughout their lives.**



### AGES 13 THROUGH 18

Once children become teenagers, there is often a shift in responsibilities over time. These young adults start taking on more of their day-to-day care themselves. After getting a driver's license,



they will likely start eating more meals away from home with their friends. It will be important to focus on a few key dietary behaviors to promote healthy kidneys over these years:

1. Avoid excess protein intake, particularly for teenagers participating in sports.
2. Eat fruits and vegetables at meal times, especially at breakfast, which many teenagers skip.
3. Watch sodium intake when eating out.
4. Drinking plenty of fluid is also important, but, hopefully, by the time they become teenagers, this has become second nature to them.

Your child may wake up last minute and choose to skip breakfast due to time constraints. **Breakfast is an opportune time for children to eat some fruit during the day, and most fruits require no preparation or cooking time.** Your child may even want to create a variety of different fruit smoothies, which will help get an assortment of fruits into the diet while taking less time to prepare than toasting a Pop-Tart® and is much healthier!

As children spend more time away from home, they are bound to eat at more fast-food restaurants. It will become increasingly important to make sure they know how to select appropriate foods from these places. It will ultimately be up to your children to make the best decisions for themselves.

If your child is involved in sports, training may become more time-consuming. Team coaches and trainers at the school may start

offering dietary guidance, especially in competitive sports, such as football and wrestling. While these coaches may have the best intentions to help your child succeed in athletics, what is good for the majority of kids may not be appropriate for your child. Often, protein intake is pushed as the most important thing to gain muscle, but high intakes of protein should be avoided in children living with PKD. It is best for parents to work with the team's coaches to help them





understand your child's medical situation. Explain that, for your child to best perform, it is best for the coach to advocate they follow the dietary guidance outlined in this cookbook. If the coaches understand what is going on and that research has led to the development of these dietary guidelines for them, then your child's coach could help advocate good dietary choices while your child engages in his or her favorite sports-related activities. When discussing diet with your child's coaches, it is best to educate them on the typical course of the disease and why it is so important for you to do everything you can to keep your child's kidneys healthy now. This will help the coaches understand the importance of not drastically changing your child's diet to help improve their sports performance, since short-term benefits in muscle growth or weight gain or loss won't outweigh the long-term benefits of making your child's kidneys last. All performance-enhancing supplements should be avoided unless approved by your child's physician.

## 18 YEARS AND OLDER

Once children reach 18 years of age, they may head off to college or consider moving out on their own. If they are going to college, keep in mind that your child may have limited access to a kitchen, equipment to cook in a dorm room or may not know how to cook. Teaching your child basic food preparation skills will be important during or even before their high-school years. Often, colleges have food passes for eating at restaurants on campus. It would be good for your child to review what his or her options are and find the most suitable places to eat to maintain the healthiest diet possible. During college, and even high school, marathon study sessions combined with limited sleep may lead your child toward drinking energy drinks or taking energy shots. It is best for these to be avoided completely, as they are loaded with caffeine and may stimulate the growth of cysts. Instead, sufficient sleep should be the goal, as well as engaging in activities that relieve stress.



# Appetizers

Mango Salsa Wontons	33
Artichoke Dip	34
Fresh Tzatziki	35
Chicken Lettuce Wraps	36
Sweet and Spicy Meatballs	38
Hot Crab Dip	39
Mexican Layer Dip	40
Hummus	42
Chili Lime Shrimp	44
Easy Bruschetta	45



**Mango Salsa Wontons**



# Mango Salsa Wontons

Servings: 24

Serving size: 1 wonton



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

## INGREDIENTS

Vegetable oil cooking spray	1 tablespoon olive oil
1 large ripe mango, peeled, pitted and diced	Pinch of cayenne pepper
1 small cucumber, peeled, seeded and diced	2 to 3 tablespoons chopped fresh cilantro (or more to taste)
½ medium red onion, finely diced	24 wonton sheets (1 package; available at supermarkets or specialty stores)
2 to 3 tablespoons fresh lime juice	

## COOKING INSTRUCTIONS

1. Preheat oven to 350° F.
2. Coat mini-muffin pans with cooking spray and line the molds with wonton sheets.
3. Bake for 9 to 12 minutes or until golden brown.
4. Set the wonton cups aside to cool.
5. Combine remaining ingredients plus pepper to taste.
6. Fill each wonton with salsa and serve.

### NUTRITION INFORMATION

(per serving)

Calories 40  
 Fat 0.7 g  
 Carbohydrates 7 g  
 Protein 1 g  
 Dietary Fiber 0.5 g  
 Calcium 4 mg  
 Phosphorus 11 mg  
 Sodium 6 mg  
 Potassium 45 mg



## APPETIZERS



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

# Artichoke Dip

Servings: 10

Serving size:  $\frac{1}{10}$  of recipe

### NUTRITION INFORMATION

(per serving)

Calories 40

Fat 2.4 g

Carbohydrates 4 g

Protein 1 g

Dietary Fiber 1.5 g

Calcium 24 mg

Phosphorus 31 mg

Sodium 113 mg

Potassium 75 mg

### INGREDIENTS

- |   |                               |
|---|-------------------------------|
| 1 cup frozen artichoke hearts               | 1 large garlic clove, crushed |
| $\frac{1}{4}$ cup low-fat mayonnaise        | 2 teaspoons hot sauce         |
| $\frac{1}{4}$ cup low-fat sour cream        | 1 tablespoon Parmesan cheese  |
| 2 tablespoons light or low-fat cream cheese |                               |

### COOKING INSTRUCTIONS

1. Preheat oven to 375° F.
2. Place artichoke hearts in a saucepan, cover with water and bring to a boil.
3. Reduce heat to medium and cook for 6 minutes.
4. Drain and rinse with cold water to cool.
5. Chop artichoke hearts.
6. In a medium bowl, combine mayonnaise, sour cream, cream cheese, hot sauce and garlic.
7. Transfer mixture to a baking dish. Top with Parmesan cheese.
8. Place in oven and bake for 30 minutes or until bubbly on top.

### NOTES

If using canned or a jar of artichokes instead of frozen, look for the lowest-sodium brand. Rinse well to help reduce sodium.

### SERVING SUGGESTION

Try this dip on toasted slices of baguette, crisp celery or with your favorite low-sodium crackers.



# Fresh Tzatziki

Servings: 10

Serving size: 2 tablespoons



LOW SODIUM



LOW PROTEIN



REDUCES ACID

## INGREDIENTS

1 large cucumber, peeled and finely chopped  
1 cup plain nonfat yogurt  
1 teaspoon dried dill weed

4 large cloves of garlic, minced  
1 tablespoon fresh lemon juice  
Pinch of salt

## COOKING INSTRUCTIONS

1. Combine all ingredients in bowl.
2. Let stand for 20 minutes.

## SERVING SUGGESTION

Serve with fresh vegetables, such as cucumbers, broccoli and carrots. Use as a dressing for salad or as a spread for a sandwich. Leftovers will keep in the refrigerator for 3 to 4 days.

### NUTRITION INFORMATION

(per serving)

Calories 20

Fat 0.1 g

Carbohydrates 3 g

Protein 2 g

Dietary Fiber 0.2 g

Calcium 57 mg

Phosphorus 47 mg

Sodium 35 mg

Potassium 110 mg



## APPETIZERS



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

# Chicken Lettuce Wraps

Servings: 6

Serving size: 2 wraps

### NUTRITION INFORMATION

(per serving)

Calories 155

Fat 5.0 g

Carbohydrates 18 g

Protein 11 g

Dietary Fiber 2.8 g

Calcium 46 mg

Phosphorus 149 mg

Sodium 364 mg

Potassium 614 mg

### INGREDIENTS

Vegetable oil cooking spray  
4 medium carrots, peeled and finely diced  
2 large (11 to 12 inches) stalks of celery, finely diced  
1 large red bell pepper, seeded and finely diced  
1 can (8 ounces) water chestnuts, drained and finely diced  
3 medium scallions (white and green parts), thinly sliced  
2 tablespoons fresh ginger, grated or finely minced  
4 cloves garlic, minced  
1 pound ground chicken (at least 90 percent lean)  
¼ teaspoon black pepper  
½ cup Chinese plum sauce  
2 tablespoons reduced-sodium soy sauce  
2 tablespoons rice vinegar  
1 teaspoon hot pepper sauce (or to taste)  
¼ cup minced fresh cilantro, plus extra for garnish  
1 head Boston or Bibb lettuce



**Dialysis-  
friendly**  
option:

This recipe contains more than 600 milligrams of potassium. For those on dialysis and watching potassium, consider limiting to 1 wrap and having a low-potassium side item with it.



**COOKING INSTRUCTIONS**

1. Liberally coat a large skillet with cooking spray and preheat over medium-high heat.
2. Add the carrots, celery, bell pepper, water chestnuts, scallions, ginger and garlic.
3. Sauté, stirring occasionally, until the vegetables soften slightly, about 5 minutes. Add a tablespoon of water as necessary to prevent scorching.
4. Reapply cooking spray, if necessary, and add the ground chicken to the skillet.
5. Cook until the chicken is no longer pink, breaking the meat into a fine crumble with a wooden spoon as it cooks.
6. Season with pepper.
7. Add the plum sauce, soy sauce, vinegar and chili paste. Stir to coat.
8. Reduce the heat to low and simmer until heated through.
9. Remove the skillet from heat and stir in cilantro.
10. Allow the mixture to cool slightly, about 5 minutes.
11. Clean the lettuce and break off 12 individual leaves.

**NOTES**

When gathering the lettuce leaves, trim away the stem end of the leaves if they are tough.

**SERVING SUGGESTION**

Fill each lettuce cup with ½ cup of the chicken mixture. Garnish with additional cilantro if desired.



## APPETIZERS



LOW  
SODIUM



LOW  
PROTEIN

# Sweet and Spicy Meatballs

Servings: 18

Serving size: 2 meatballs

### NUTRITION INFORMATION

(per serving)

Calories 80

Fat 2.9 g

Carbohydrates 8 g

Protein 5 g

Dietary Fiber 0.4 g

Calcium 11 mg

Phosphorus 43 mg

Sodium 34 mg

Potassium 81 mg

### INGREDIENTS

Vegetable oil cooking spray  
¼ cup onion, chopped  
1 pound 85 percent or leaner  
ground beef  
½ cup fine dry breadcrumbs  
¼ cup fresh parsley, chopped  
⅛ teaspoon nutmeg  
¼ cup liquid reduced-fat or  
low-fat nondairy creamer

1 large egg white, beaten  
½ cup fresh or frozen  
cranberries, finely chopped  
2 teaspoons dry mustard  
⅛ teaspoon cayenne pepper  
½ cup grape jelly  
1 teaspoon fresh lemon juice

### COOKING INSTRUCTIONS

1. Preheat oven to 375° F.
2. Coat a small saucepan with cooking spray and place over medium heat.
3. Add onion and sauté until tender.
4. Combine onion with next 6 ingredients in a bowl.
5. Shape into 36 1-inch meatballs.
6. Coat a jelly roll pan or shallow baking dish with cooking spray.
7. Place meatballs in pan and bake for 18 minutes.
8. Meanwhile, prepare sauce by combining the cranberries and remaining ingredients in a small saucepan.
9. Cook over medium heat until heated through.
10. Transfer meatballs to a serving dish and pour sauce over meatballs. Serve with toothpicks.

### SERVING SUGGESTION

Pair with some vegetables and dip to reduce the dietary acids, and you have the makings of a PKD-friendly appetizer!





# Hot Crab Dip

Servings: 10

Serving size: 1/10 of recipe



LOW SODIUM



LOW PROTEIN

## INGREDIENTS

- |  |                                     |
|--|-------------------------------------|
| 1 package (8 ounces) light or low-fat cream cheese, softened | 1/8 teaspoon black pepper           |
| 1 tablespoon onion, finely minced                            | Cayenne pepper to taste             |
| 1 teaspoon fresh lemon juice                                 | 2 tablespoons nondairy creamer      |
| 2 teaspoons low-sodium Worcestershire sauce                  | 1 can (6 ounces) crab meat, drained |

## COOKING INSTRUCTIONS

1. Preheat oven to 375° F.
2. Place softened cream cheese in a bowl.
3. Add onion, lemon juice, Worcestershire sauce, black pepper and cayenne pepper. Mix well.
4. Stir in nondairy creamer.
5. Add crab meat and stir until blended.
6. Place mixture into an oven-safe dish.
7. Bake uncovered for 15 minutes or until hot and bubbly.

## SERVING SUGGESTION

Serve warm with low-sodium crackers.

### NUTRITION INFORMATION

(per serving)

- Calories 65
- Fat 3.7 g
- Carbohydrates 3 g
- Protein 5 g
- Dietary Fiber 0 g
- Calcium 51 mg
- Phosphorus 78 mg
- Sodium 179 mg
- Potassium 117 mg



## APPETIZERS



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

### NUTRITION INFORMATION

(per serving)

Calories 110

Fat 6.7 g

Carbohydrates 11 g

Protein 4 g

Dietary Fiber 4.2 g

Calcium 53 mg

Phosphorus 88 mg

Sodium 66 mg

Potassium 347 mg

# Mexican Layer Dip

Servings: 15

Serving size: about 4 bites

## INGREDIENTS

- 1 15-ounce can low-sodium refried black beans
- 3 ripe avocados, peeled and pitted
- 1 tablespoon fresh lemon juice
- 1 tablespoon light mayonnaise
- ½ cup low-fat sour cream
- 2 tablespoons fresh cilantro, chopped
- 3 tablespoons low-sodium taco seasoning
- 2 medium bell peppers, any color, diced
- ½ cup of your favorite low-sodium cheese, shredded
- 2 to 3 cups shredded lettuce
- ½ cup diced plum tomato
- ½ cup diced onion



## **COOKING INSTRUCTIONS**

1. Spread ingredients in 9" x 11" baking pan.
2. Layer 1: refried black beans
3. Layer 2: Mash together: 3 ripe avocados, lemon juice and mayonnaise
4. Layer 3: Mix sour cream, cilantro and low-sodium taco seasoning
5. Layer 4: bell peppers
6. Layer 5: shredded cheese
7. Layer 6: shredded lettuce
8. Layer 7: diced plum tomato and onion

## **NOTES**

Substitute Homemade Salt-Free Mexican Spice Mix on page 103 for the low-sodium taco seasoning, and make your own light mayonnaise with the recipe on page 104.

For low-sodium cheese, we suggest Monterey jack or low-sodium versions of cheddar or pepper jack. You can also mix a blend for variety.

## **SERVING SUGGESTIONS**

The joy in this dish is how much you can personalize it to your taste. If you don't like something, just leave it out. Is it missing your favorite taco ingredient? Add it in and make it your own.

Serve with raw vegetables such as baby carrots, broccoli florets and cauliflower florets, salt-free tortilla chips or tortillas cut into triangles.



## APPETIZERS



LOW  
SODIUM



LOW  
PROTEIN

# Hummus

Servings: 8

Serving size:  $\frac{1}{8}$  of recipe

### NUTRITION INFORMATION

(per serving)

Calories 95

Fat 3.9 g

Carbohydrates 12 g

Protein 4 g

Dietary Fiber 3.3 g

Calcium 47 mg

Phosphorus 104 mg

Sodium 28 mg

Potassium 157 mg

### INGREDIENTS

$\frac{3}{8}$ cup dry or $1\frac{1}{2}$ cups cooked and drained garbanzo beans (chickpeas)	2 to 3 tablespoons tahini sauce
Juice from 1 lemon	2 tablespoons onion, minced
2 cloves garlic, minced	2 tablespoons fresh parsley, minced
	Pinch of salt

### COOKING INSTRUCTIONS

#### Beans:

1. Place beans in a large bowl and cover with several inches of cold water. They will double in size during soaking, so be sure to use a large enough bowl to accommodate.
2. Remove and discard any floating beans.
3. Cover the bowl with a clean towel and let soak overnight. Drain and rinse before cooking. Place beans in large pot and cover with water (about 1 quart of water for each cup of soaked beans).
4. Bring to a boil and reduce to a simmer.
5. Cook for 60 to 90 minutes until tender. Watch water level as you cook, you may need to add more water.
6. Drain in colander and allow to cool.
7. Place cooked and drained beans in airtight container or sealable bag (keep liquid out). They will keep in the refrigerator for 3 to 4 days.

#### Hummus:

1. Mash cooked garbanzo beans into thick paste.
2. Add the rest of the ingredients and stir together.
3. If consistency is too thick, add a bit more lemon juice.



**NOTES**

Using dried garbanzo beans, rather than canned, allows you to better control the salt content. Consider cooking a large batch and freezing the extras for use next time. To freeze, pat dry with a paper towel before adding to a sealable bag. Frozen garbanzos will keep for about a year.

Hummus requires quite tender beans, but they freeze better when they are firmer. You may want to remove a portion to freeze and continue cooking the rest.

**SERVING SUGGESTION**

Serve with fresh pita bread and fresh vegetables such as cucumbers, broccoli and carrots for a snack.



LOW  
SODIUM



HIGH  
ACID

# Chili Lime Shrimp

Servings: 4

Serving size: about 5 shrimp

## INGREDIENTS

¼ cup low-sodium or no-salt-added tomato paste	1 pound fresh medium shrimp (16 to 20 count), peeled, deveined
1 clove garlic, minced	
1 teaspoon chili powder	1 lime, cut into wedges
Vegetable oil cooking spray	

## NUTRITION INFORMATION

(per serving)

Calories 95

Fat 0.8 g

Carbohydrates 2 g

Protein 20 g

Dietary Fiber 0.4 g

Calcium 67 mg

Phosphorus 203 mg

Sodium 105 mg

Potassium 271 mg

## COOKING INSTRUCTIONS

1. Combine tomato paste, garlic and chili powder in small bowl.
2. Spray wok or large skillet with cooking spray; heat on high 1 minute.
3. Add shrimp, stirring frequently; cook 5 minutes or until shrimp turn pink.
4. Add tomato paste mixture; toss to evenly coat shrimp.

## NOTES

Use fresh shrimp, as frozen are often high in sodium.

## SERVING SUGGESTIONS

Divide shrimp evenly among 4 serving plates. Squeeze 1 lime wedge over shrimp on each plate before serving. Serve with additional lime wedges.

To reduce the acids from the meat, pair with some vegetable sides or fresh fruit.



# Easy Bruschetta

Servings: 8

Serving size: 6 slices



LOW SODIUM



LOW PROTEIN

## INGREDIENTS

½ cup red bell pepper, chopped	1½ teaspoons dried basil leaves
½ cup dry-packed, sun-dried tomatoes, chopped	1½ teaspoons dried oregano leaves
2 tablespoons grated Romano cheese	1 baguette about 26 inches long, cut into ½-inch diagonal slices, toasted
2 tablespoons grated Parmesan cheese	1 can (14½ ounces) low-sodium or no-salt-added petite diced tomatoes
1 tablespoon minced garlic	

## COOKING INSTRUCTIONS

1. Combine and mix all ingredients, except baguette slices, in a saucepan.
2. Season with pepper to taste; bring to a boil.
3. Reduce heat and let simmer for 10 minutes.

## SERVING SUGGESTION

Serve at room temperature on toasted baguette slices.

### NUTRITION INFORMATION


(per serving)

Calories 170  
 Fat 1.9 g  
 Carbohydrates 31 g  
 Protein 8 g  
 Dietary Fiber 2.1 g  
 Calcium 78 mg  
 Phosphorus 99 mg  
 Sodium 368 mg  
 Potassium 265 mg



# Soups and Salads

Eggplant Vegetable Soup	47
Turkey Vegetable Soup	48
Ginger Pork Soup	49
Creamy Carrot Thyme Soup	50
Beef Chili	51
Strawberry Spinach Salad	53
Roasted Corn and Edamame Salad	54
Summertime Salad	55
Zucchini-Ribbon Salad	56
Mandarin Salad	57
Chipotle Chicken Salad	58
Spinach, Apple and Walnut Salad	59
Mexican Vegetable Salad	60
Fall Harvest Orzo Salad	61



**Eggplant Vegetable Soup**





# Eggplant Vegetable Soup

Servings: 12

Serving size: 1 cup



LOW  
SODIUM



HIGH  
FLUID

## INGREDIENTS

1 pound 93 percent lean ground turkey	1 medium eggplant, peeled and cubed
½ cup onion, chopped	1 clove garlic, minced
½ cup celery, chopped	½ teaspoon salt
½ cup carrots, chopped	1½ teaspoon ground nutmeg
1 can (28 ounces) low-sodium or no-salt-added crushed tomatoes with liquid	½ cup dry macaroni
28 ounces low-sodium or no-salt-added beef broth	2 teaspoons dried parsley
	Pepper to taste
	¾ cup grated Parmesan cheese for garnish

## NUTRITION INFORMATION

(per serving)

**Calories** 145  
**Fat** 5.6 g  
**Carbohydrates** 12 g  
**Protein** 13 g  
**Dietary Fiber** 2.2 g  
**Calcium** 106 mg  
**Phosphorus** 156 mg  
**Sodium** 252 mg  
**Potassium** 312 mg

## COOKING INSTRUCTIONS

1. In a pot, brown turkey.
2. Add onions, celery, carrots, tomatoes and broth.
3. Stir well.
4. Add eggplant and spices; simmer for 30 minutes.
5. Add macaroni, cook 10 minutes longer, until macaroni is done.
6. Add parsley.
7. Serve sprinkled with cheese.

## SERVING SUGGESTION

This hearty soup is loaded with healthy vegetables. Want it as a side dish instead? Considering making it without the meat.





# Turkey Vegetable Soup

Servings: 12

Serving size: 1 cup

## NUTRITION INFORMATION

(per serving)

Calories 245  
 Fat 13.8 g  
 Carbohydrates 8 g  
 Protein 24 g  
 Dietary Fiber 1.5 g  
 Calcium 99 mg  
 Phosphorus 270 mg  
 Sodium 125 mg  
 Potassium 498 mg

## INGREDIENTS

¼ cup unsalted butter	2 tablespoons fresh parsley, chopped
2 medium onions, chopped	½ teaspoon fresh sage, chopped
2 tablespoons flour	2 pounds cooked 93 percent lean ground turkey
1½ teaspoon low-sodium curry powder	1½ cups milk
3 cups low-sodium chicken broth	10 ounces frozen chopped spinach
1 cup potatoes, chopped	Pepper to taste
½ cup celery, chopped	
½ cup carrots, chopped	

## COOKING INSTRUCTIONS

1. Melt butter in pot over medium-high heat.
2. Add onions and sauté until translucent, approximately 10 minutes.
3. Stir in flour and curry powder, cook 2 to 3 minutes.
4. Add broth, potatoes, carrots, celery, parsley and sage.
5. Bring to a boil.
6. Reduce heat to low, cover and simmer for 10 minutes.
7. Add turkey, milk and spinach.
8. Cover and simmer until heated through.
9. Add pepper to taste.

## SERVING SUGGESTION

Having this soup, along with a small salad topped with your favorite vegetables, is a vegetable medley that'll certainly help protect your kidneys.



# Ginger Pork Soup

Servings: 8

Serving size: 1 cup



## INGREDIENTS

1 tablespoon olive oil	2 tablespoons dry sherry
12 ounces lean boneless pork, cut into thin pieces	2 tablespoons low-sodium soy sauce
2 cups shiitake mushrooms, sliced	2 teaspoons fresh ginger
2 cloves garlic, minced	½ teaspoon crushed red pepper
32 ounces low-sodium chicken broth or see Basic Chicken Stock recipe on page 108	2 cups Chinese cabbage, thinly sliced
	1 scallion, thinly sliced

## COOKING INSTRUCTIONS

1. Add oil to large saucepan, warm on medium heat.
2. Add pork and cook for 2 to 3 minutes or until slightly pink in the center.
3. Remove pork from pan and set aside.
4. Add mushrooms and garlic to saucepan and cook until tender.
5. Stir in broth, sherry, soy sauce, ginger and red pepper and bring to a boil.
6. Stir in pork, Chinese cabbage and green onion.
7. Heat until hot throughout.

## SERVING SUGGESTION

Try pairing with a side salad for a light dinner.

### NUTRITION INFORMATION

(per serving)

<b>Calories</b>	140
<b>Fat</b>	6.0 g
<b>Carbohydrates</b>	8 g
<b>Protein</b>	13 g
<b>Dietary Fiber</b>	1.1 g
<b>Calcium</b>	28 mg
<b>Phosphorus</b>	134 mg
<b>Sodium</b>	194 mg
<b>Potassium</b>	356 mg





LOW SODIUM



LOW PROTEIN



HIGH FLUID



REDUCES ACID

# Creamy Carrot Thyme Soup

Servings: 12

Serving size: 1 cup

## NUTRITION INFORMATION

(per serving)

Calories 100

Fat 2.6 g

Carbohydrates 17 g

Protein 3 g

Dietary Fiber 3.2 g

Calcium 50 mg

Phosphorus 65 mg

Sodium 433 mg

Potassium 393 mg

## INGREDIENTS

- |  |                          |
|--|--------------------------|
| 3 pounds baby carrots or carrot chunks, peeled                         | 2 sprigs fresh thyme     |
| 8 cups low-sodium vegetable or chicken broth (see page 108 for recipe) | ½ teaspoon ground ginger |
|  | ⅓ cup honey              |
|  | ⅓ cup heavy cream        |
|  | Pepper to taste          |

## COOKING INSTRUCTIONS

1. Combine the carrots, stock and thyme in a pot.
2. Bring to a boil, reduce the heat to low and simmer for 45 minutes to 1 hour.
3. Remove from heat and transfer the soup in 2 batches to a blender.
4. Puree the soup completely.
5. Stir in the honey and cream.
6. Add pepper, as desired.
7. Serve while warm.



# Beef Chili

Servings: 8

Serving size: 1 cup



LOW  
SODIUM



HIGH  
ACID

## INGREDIENTS

½ cup onion, chopped	16 ounces low-sodium stewed
1 large (11 to 12 inches) stalk of celery	tomatoes
½ cup green bell pepper, chopped	1 tablespoon canola oil
1½ pounds 85 percent or leaner ground beef	2 tablespoons chili powder
	1½ cup water

## COOKING INSTRUCTIONS

1. Chop onion, celery and bell pepper.
2. Heat large skillet on medium heat. Add oil, onion, celery and bell pepper. Cook until tender but not brown.
3. Add ground beef, breaking into small pieces, and cook until brown.
4. Blend tomatoes in a blender and add to beef mixture.
5. Add chili powder and water. Mix thoroughly; reduce heat to low.
6. Simmer for several hours.

## SERVING SUGGESTION

This beef chili is far healthier than the canned products you find at the store. Serve with some roasted vegetables or mixed vegetables to top off the meal.

### NUTRITION INFORMATION

(per serving)

Calories 175

Fat 10.5 g

Carbohydrates 5 g

Protein 16 g

Dietary Fiber 1.4 g

Calcium 37 mg

Phosphorus 130 mg

Sodium 83 mg

Potassium 330 mg





# Strawberry Spinach Salad

Servings: 6

Serving size: 1/6 of recipe



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

## INGREDIENTS

3 cups fresh spinach or 6-ounce package of pre-washed spinach	1 tablespoon red wine vinegar
1 cup fresh strawberries, finely sliced	1 tablespoon fresh lemon juice
	2 tablespoons honey
	1/8 teaspoon dry mustard
	3 tablespoons olive oil

## COOKING INSTRUCTIONS

1. Wash and dry the spinach leaves.
2. Tear into bite-size pieces and place in a bowl.
3. Top spinach with sliced strawberries.
4. In a separate bowl, mix together the vinegar, lemon juice, honey, dry mustard and oil.
5. Pour dressing over spinach and strawberries.
6. Toss well and serve immediately.

### NUTRITION INFORMATION

(per serving)

Calories 95

Fat 6.9 g

Carbohydrates 9 g

Protein 1 g

Dietary Fiber 0.9 g

Calcium 20 mg

Phosphorus 15 mg

Sodium 13 mg

Potassium 134 mg



LOW  
SODIUMLOW  
PROTEINREDUCES  
ACID

# Roasted Corn and Edamame Salad

Servings: 6

Serving size: 1/6 of recipe

## NUTRITION INFORMATION

(per serving)

Calories 65

Fat 2.0 g

Carbohydrates 10 g

Protein 3 g

Dietary Fiber 1.6 g

Calcium 26 mg

Phosphorus 52 mg

Sodium 71 mg

Potassium 178 mg

## INGREDIENTS

- |                                 |                                |
|---------------------------------|--------------------------------|
| 2 ears fresh corn, unhusked, or | 1 tablespoon light mayonnaise  |
| 1 1/4 cups cooked corn kernels  | 1 tablespoon fresh lemon juice |
| 1/2 cup edamame, shelled        | 1 1/2 teaspoon ginger, finely  |
| 1/4 cup red onion, chopped      | chopped or grated              |
| 1/4 cup red bell pepper, diced  | 1/8 teaspoon salt              |
| small                           | 1/8 teaspoon freshly ground    |
| 1 tablespoon fresh cilantro,    | black pepper                   |
| finely chopped                  |                                |

## COOKING INSTRUCTIONS

1. Soak fresh corn in cold water about 30 minutes.
2. Heat grill on high.
3. Grill corn in husk, 10 to 15 minutes, turning once.
4. Let cool and remove husks.
5. Cut corn from cob into a bowl.
6. Combine with remaining ingredients.
7. Cover and chill in refrigerator until ready to serve.

## NOTES

Edamame should be cooked prior to chilling in the refrigerator. Frozen varieties may be pre-cooked prior to packaging. Check package for details.





# Summertime Salad

Servings: 6      Serving size: 1½ cups

Recipe submitted by Katelynne Burghart



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

## INGREDIENTS

### Salad:

6 cups spring lettuce mix, loosely packed	½ cup goat cheese, crumbled
2 cups seedless grapes, halved	½ cup walnuts, toasted and chopped
1 avocado, peeled, pitted and diced	¼ cup red onion, thinly sliced

### Creamy Poppy Seed dressing:

3 tablespoons apple cider vinegar	⅛ teaspoon salt
¼ cup sugar	3 tablespoons light mayonnaise
1 tablespoon onion, finely chopped	1 tablespoon freshly squeezed orange juice
⅛ teaspoon dry mustard	3 tablespoons vegetable oil
	½ teaspoon poppy seeds

### NUTRITION INFORMATION

(per serving)

Calories 305

Fat 22.1 g

Carbohydrates 24 g

Protein 6 g

Dietary Fiber 3.9 g

Calcium 85 mg

Phosphorus 123 mg

Sodium 185 mg

Potassium 457 mg

## COOKING INSTRUCTIONS

1. Combine vinegar, sugar, onion, dry mustard, salt, mayonnaise and orange juice in a food processor.
2. Process to blend well.
3. With processor still on, add vegetable oil in a steady stream and continue processing until incorporated.
4. Add poppy seeds and pulse briefly until just blended.
5. Transfer to airtight container and refrigerate until ready to use.
6. Add salad ingredients to a bowl.
7. Drizzle with vinaigrette, toss to combine.





LOW SODIUM



LOW PROTEIN



REDUCES ACID

# Zucchini-Ribbon Salad

Servings: 6

Serving size:  $\frac{1}{6}$  of recipe

## INGREDIENTS

2 medium zucchini

2 medium yellow squash

3 tablespoons low-sodium soy sauce

2 tablespoons rice wine vinegar

1 tablespoon sesame oil

$\frac{1}{4}$  teaspoon sugar

$\frac{1}{8}$  teaspoon red pepper flakes

$\frac{1}{2}$  teaspoon sesame seeds

## NUTRITION INFORMATION

(per serving)

Calories 50

Fat 2.8 g

Carbohydrates 5 g

Protein 2 g

Dietary Fiber 1.5 g

Calcium 22 mg

Phosphorus 60 mg

Sodium 272 mg

Potassium 358 mg

## COOKING INSTRUCTIONS

1. Peel long ribbons of the zucchini and squash.
2. In a large bowl, whisk together the remaining ingredients.
3. Add the zucchini and squash, and toss to mix well.
4. Season to taste with freshly ground pepper.



# Mandarin Salad

Servings: 4

Serving size: ¼ of recipe



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

## INGREDIENTS

### Salad:

4 cups leaf lettuce or spinach	¼ cup toasted almonds
1 cup mandarin oranges, packed in water or juice	3 to 5 strips low-sodium bacon or turkey bacon, cooked and crumbled
½ cup dried cherries	
2 medium apples, sliced	

### Dressing:

2 tablespoons sugar substitute (Splenda, or can use sugar)	¼ teaspoon salt
2 tablespoons apple cider vinegar	2 tablespoons water
	2 tablespoons olive oil
	Pepper to taste

### NUTRITION INFORMATION

(per serving)

Calories 295

Fat 14.8 g

Carbohydrates 37 g

Protein 7 g

Dietary Fiber 5.2 g

Calcium 67 mg

Phosphorus 121 mg

Sodium 245 mg

Potassium 476 mg

## COOKING INSTRUCTIONS

1. Mix salad ingredients.
2. In a separate bowl, mix ingredients for dressing in the order they are listed.
3. Toss salad with dressing just before serving.





LOW  
SODIUM



HIGH  
ACID

# Chipotle Chicken Salad

Servings: 4

Serving size: ¼ of recipe

Recipe submitted by Emily Decker

## INGREDIENTS

2 cups cooked chicken, shredded	2 tablespoons chipotle peppers in adobo sauce
1 cup celery, chopped	¼ teaspoon sea salt
¼ cup red onions, diced	⅛ teaspoon black pepper
½ cup light or homemade mayonnaise (see page 104 for recipe)	

### NUTRITION INFORMATION

(per serving)

Calories 195

Fat 9.8 g

Carbohydrates 5 g

Protein 21 g

Dietary Fiber 0.6 g

Calcium 25 mg

Phosphorus 159 mg

Sodium 442 mg

Potassium 258 mg

## COOKING INSTRUCTIONS

1. Combine all ingredients in large bowl.
2. Mix well.

## SERVING SUGGESTION

Serve on lettuce leaves, low-salt crackers or bread roll. Don't forget to pair with a side of mixed fruit or grapes to make sure you keep your dietary acid load low.



# Spinach, Apple and Walnut Salad

Servings: 4

Serving size: ¼ of recipe



LOW  
SODIUM



LOW  
PROTEIN



HIGH  
FLUID



REDUCES  
ACID

## INGREDIENTS

2 large apples (Golden Delicious or other favorite), cored, diced large	1 tablespoon apple cider vinegar
¼ cup fresh lemon juice	2 tablespoons honey
8 cups (2 5-ounce packages) baby spinach leaves	Ground black pepper
3 tablespoons extra-virgin olive oil	⅔ cup crumbled goat cheese
	½ cup chopped walnuts, toasted

## COOKING INSTRUCTIONS

1. Toss apples with 2 tablespoons of the lemon juice.
2. Place spinach in a large bowl; remove long stems and bruised leaves.
3. Whisk together remaining juice, olive oil, vinegar, honey and ground pepper to taste.
4. Toss spinach with apples and dressing.

## SERVING SUGGESTION

Divide among 4 bowls. Top with cheese and walnuts.

### NUTRITION INFORMATION

(per serving)

<b>Calories</b>	380
<b>Fat</b>	27.1 g
<b>Carbohydrates</b>	29 g
<b>Protein</b>	9 g
<b>Dietary Fiber</b>	5.1 g
<b>Calcium</b>	153 mg
<b>Phosphorus</b>	182 mg
<b>Sodium</b>	170 mg
<b>Potassium</b>	580 mg



**Dialysis-friendly**  
option:

Due to potassium content, individuals on dialysis and watching their potassium intake should either make this salad an entire meal or select another salad lower in potassium.





LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

# Mexican Vegetable Salad

Servings: 6

Serving size: 1/6 of recipe

Recipe submitted by Nicole Harr

## NUTRITION INFORMATION

(per serving)

Calories 70

Fat 4.8 g

Carbohydrates 7 g

Protein 1 g

Dietary Fiber 1.9 g

Calcium 24 mg

Phosphorus 34 mg

Sodium 88 mg

Potassium 288 mg

## INGREDIENTS

- |   |                                       |
|---|---------------------------------------|
| 2 medium tomatoes, seeded and chopped               | 1 jalapeño, seeded and finely chopped |
| 1 cucumber, cut into bite-size chunks               | 2 tablespoons fresh cilantro          |
| 1/2 medium red onion, chopped                       | 2 teaspoons Tabasco                   |
| 1 medium red bell pepper, cut into chunks           | 2 limes, juiced                       |
| 2 large (11 to 12 inches) stalks of celery, chopped | 2 tablespoons olive oil               |
|   | Salt and pepper to taste              |

## COOKING INSTRUCTIONS

1. Combine vegetables in a bowl.
2. Sprinkle with chopped cilantro.
3. Dress salad with hot sauce, lime juice and olive oil.
4. Season with salt and pepper.
5. Serve chilled.



# Fall Harvest Orzo Salad

Servings: 8

Serving size: 1¼ cup



LOW  
SODIUM



LOW  
PROTEIN

## INGREDIENTS

4 cups cooked orzo, chilled (about 1⅔ cup dried orzo)	2 tablespoons fresh basil, chopped
1 cup dried cranberries	½ cup crumbled blue cheese
2 cups apple, diced	¼ cup blanched almonds, chopped
¼ cup extra-virgin olive oil	
¼ cup fresh lemon juice	
½ teaspoon freshly ground black pepper	

## COOKING INSTRUCTIONS

1. Prepare the orzo as directed on the packaging.
2. In a medium bowl, add all of the ingredients except blue cheese and almonds, gently combining until well incorporated.
3. Transfer the mixture to a serving dish, sprinkle with the crumbled blue cheese and almonds and serve.

### NUTRITION INFORMATION

(per serving)

**Calories** 305

**Fat** 13.5 g

**Carbohydrates** 40 g

**Protein** 7 g

**Dietary Fiber** 3.6 g

**Calcium** 71 mg

**Phosphorus** 110 mg

**Sodium** 121 mg

**Potassium** 146 mg



# Sides

Warm Potato and Kale Mix	63
Spicy Sweet Potato Fries	64
Mashed Tomato Potatoes	65
Grilled Sweet Potatoes and Scallions	66
Roasted Brussels Sprouts	67
Grilled Asparagus With Mozzarella	68
Grilled Vegetables	69
Noodles Romano	70
Quinoa With Black Beans and Avocado	71
Roasted Vegetables	72
Couscous and Feta-Stuffed Peppers	73



Warm Potato  
and Kale Mix





# Warm Potato and Kale Mix

Servings: 2

Serving size: ½ of recipe



LOW SODIUM



LOW PROTEIN



HIGH FLUID



REDUCES ACID

## INGREDIENTS

2 large potatoes, peeled	½ teaspoon dried thyme
1 cup fresh kale leaves, packed	1 small tomato, chopped
1 tablespoon olive oil	½ cup low-sodium white beans, drained and rinsed
½ cup onion, chopped	Pepper to taste
1 clove garlic, minced	

## COOKING INSTRUCTIONS

1. Dice potatoes and steam until just tender (about 10 minutes). Set aside.
2. Wash and remove stems from kale, then chop into ½-inch slices.
3. In nonstick skillet, heat oil on medium.
4. Add onion, garlic and thyme; sauté until onion is soft (about 3 minutes).
5. Add kale and tomato, sauté for an additional 1 to 2 minutes or until kale has wilted.
6. Add potatoes and beans, stir until mixed and warm throughout.
7. Season with pepper to taste and serve immediately.

### NUTRITION INFORMATION

(per serving)

Calories	350
Fat	7.4 g
Carbohydrates	65 g
Protein	10 g
Dietary Fiber	9.1 g
Calcium	104 mg
Phosphorus	178 mg
Sodium	162 mg
Potassium	1,233 mg



**Dialysis-friendly**  
option:

For individuals on dialysis and watching potassium intake, this recipe is higher in potassium, even with boiling potatoes, and should be avoided.





LOW SODIUM



LOW PROTEIN



REDUCES ACID

# Spicy Sweet Potato Fries

Servings: 6

Serving size: 1 potato

Recipe submitted by Nicole Harr

## INGREDIENTS

6 medium sweet potatoes (about 2¼ pounds)	½ teaspoon salt (optional)
Vegetable cooking spray	¼ teaspoon ground red pepper
2 teaspoons sugar	⅛ teaspoon black pepper

### NUTRITION INFORMATION

(per serving)

Calories 115

Fat 0.8 g

Carbohydrates 25 g

Protein 2 g

Dietary Fiber 3.8 g

Calcium 44 mg

Phosphorus 62 mg

Sodium 238 mg

Potassium 544 mg

## COOKING INSTRUCTIONS

1. Preheat oven to 500° F.
2. Peel potatoes; cut lengthwise.
3. Place in a large bowl; coat with cooking spray.
4. Combine sugar, salt and peppers and sprinkle over potatoes, tossing well to coat.
5. Arrange potatoes, cut sides down, in a single layer on a baking sheet.
6. Bake for 10 minutes; turn wedges over.
7. Bake an additional 10 minutes or until tender and beginning to brown.



**Dialysis-friendly**  
option:

Individuals on dialysis and watching their potassium intake should boil potatoes prior to baking (see page 19). Doing this will reduce the potassium by 50 percent. If you cut the fries thinner, more potassium can be removed.



# Mashed Tomato Potatoes

Servings: 8

Serving size: 1/8 of recipe



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

## INGREDIENTS

2 pounds Yukon Gold potatoes (about 6), peeled and cut into 2-inch pieces	3 medium scallions, trimmed and chopped
2 garlic cloves, peeled	1/2 cup grated Parmesan cheese
1/2 cup milk	1/2 teaspoon kosher salt
1/4 cup unsalted butter	5 medium tomatoes, chopped
1/4 cup fresh flat-leaf parsley, chopped	

## COOKING INSTRUCTIONS

1. Place the potatoes and garlic in a large saucepan and cover with water.
2. Bring to a boil; cover and simmer about 20 minutes or until a fork easily pierces a potato.
3. Drain the potatoes and garlic.
4. Mash with a potato masher or a fork until smooth.
5. Blend in the milk, butter, parsley, scallions, Parmesan and salt.
6. Gently fold in the tomatoes.

### NUTRITION INFORMATION

(per serving)

**Calories** 190  
**Fat** 8.1 g  
**Carbohydrates** 25 g  
**Protein** 6 g  
**Dietary Fiber** 3.2 g  
**Calcium** 114 mg  
**Phosphorus** 125 mg  
**Sodium** 262 mg  
**Potassium** 578 mg



**Dialysis-  
friendly**  
option:

For individuals on dialysis and watching potassium intake, this recipe is higher in potassium, even with boiling potatoes, and should be avoided.





LOW SODIUM



LOW PROTEIN



HIGH FLUID



REDUCES ACID

# Grilled Sweet Potatoes and Scallions

Servings: 4

Serving size: ¼ of recipe

## NUTRITION INFORMATION

(per serving)

Calories 570

Fat 41.3 g

Carbohydrates 46 g

Protein 5 g

Dietary Fiber 7.1 g

Calcium 106 mg

Phosphorus 123 mg

Sodium 257 mg

Potassium 1,007 mg

## INGREDIENTS

4 large sweet potatoes,  
par-cooked and cut into  
½-inch slices

8 medium scallions, sliced

¾ cup olive oil, divided

2 tablespoons Dijon mustard

½ cup apple cider vinegar

¼ cup balsamic vinegar

2 teaspoons honey

Freshly ground pepper

¼ cup fresh flat-leaf parsley,  
coarsely chopped

## COOKING INSTRUCTIONS

1. Preheat grill to high.
2. Brush potatoes and scallions with oil, and arrange on grill.
3. Grill potatoes for 3 to 4 minutes on each side, or until just tender.
4. Grill scallions until softened and marked.
5. Remove scallions from the grill and cut into thin slices.
6. In a large bowl, whisk together ½ cup olive oil, mustard, vinegars and honey. Season with pepper to taste.
7. Add potatoes, scallions and parsley, and toss until potatoes are well coated.
8. Transfer to a platter and serve.



**Dialysis-friendly**  
option:

For individuals on dialysis and watching potassium intake, avoid this recipe due to the potassium content. If you want a potato, consider using the methods discussed on page 19.



# Roasted Brussels Sprouts

Servings: 6      Serving size: ½ cup

Recipe submitted by Nicole Harr



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

## INGREDIENTS

1 pound Brussels sprouts, rinsed, ends trimmed, rough outer leaves of larger sprouts removed	1 teaspoon fresh lemon juice
1 tablespoon garlic, minced (about 3 cloves)	2 tablespoons olive oil
	Salt
	Freshly ground black pepper
	¼ cup Parmesan cheese, freshly grated (optional)

## COOKING INSTRUCTIONS

1. Preheat oven to 350° F.
2. Place Brussels sprouts in a cast-iron frying pan or a roasting pan.
3. Toss in the garlic.
4. Sprinkle Brussels sprouts with lemon juice.
5. Toss with oil so that the sprouts are well coated.
6. Sprinkle generously with salt (at least a ½ teaspoon) and a few turns of black pepper mill.
7. Put Brussels sprouts in oven on top rack, cook for 20 minutes, then stir to coat sprouts with the oil in the pan.
8. Cook for another 10 minutes.
9. Sprinkle with Parmesan (if using) and cook for another 5 minutes. Serve warm.

## NOTES

Adjust the timing depending on the size of the sprouts. They should be nicely browned, some of the outside leaves crunchy, and the interior should be cooked through. If the sprouts look like they are getting too brown (they should be well browned, not burned), move them to a lower rack or reduce the heat.

### NUTRITION INFORMATION

(per serving)

Calories 85

Fat 6.1 g

Carbohydrates 6 g

Protein 4 g

Dietary Fiber 2.0 g

Calcium 76 mg

Phosphorus 74 mg

Sodium 128 mg

Potassium 244 mg





# Grilled Asparagus With Mozzarella

Servings: 4

Serving size: 5 asparagus stalks

## INGREDIENTS

### Asparagus:

20 large asparagus stalks,  
peeled

¼ pound fresh part-skim or  
low-fat mozzarella, cut into  
4 half-inch thick slices

### Dressing:

4 tablespoons fresh lemon juice  
1 small shallot, finely chopped  
1 tablespoon fresh parsley,  
finely chopped

1 teaspoon dried oregano  
¼ cup olive oil  
Freshly ground pepper to taste

### NUTRITION INFORMATION

(per serving)

Calories 230  
Fat 19.5 g  
Carbohydrates 7 g  
Protein 10 g  
Dietary Fiber 2.2 g  
Calcium 239 mg  
Phosphorus 203 mg  
Sodium 200 mg  
Potassium 274 mg

## COOKING INSTRUCTIONS

1. Preheat grill.
2. In small bowl, mix all of the dressing ingredients together and set aside.
3. Brush asparagus with olive oil mixture, and season with pepper to taste.
4. Grill for 3 to 4 minutes or until just tender.
5. Divide the asparagus among 4 plates and immediately top with a slice of mozzarella.



**Dialysis-  
friendly**  
option:

Individuals on dialysis should reduce cheese by half to reduce the phosphorus content of this recipe. Also, remember to take your phosphorus binders if they have been prescribed to you.



# Grilled Vegetables

Servings: 4

Serving size: ¼ of recipe



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

## INGREDIENTS

- |   |  |
|---|--|
| 1 medium onion, cut into large chunks         | 1 small sweet potato, cut into small cubes |
| 1 medium zucchini, cut into thick slices      | 2 tablespoons olive oil                    |
| 1 medium yellow squash, cut into thick slices | 1 clove garlic, minced                     |
|   | Pinch of salt                              |
|   | Pepper to taste                            |

## COOKING INSTRUCTIONS

1. Put all vegetables in large bowl.
2. Mix oil, garlic, salt and pepper in small bowl and pour over vegetables. Stir until all vegetables are coated.
3. Arrange vegetables on flat baking sheet or grill basket. Cook on medium-high grill until vegetables are cooked and just starting to brown, approximately 10 to 20 minutes.
4. Serve immediately.

## NOTES

Cover the baking sheet with foil for easy cleaning. Make it your own by using any vegetables you like.

### NUTRITION INFORMATION

(per serving)

Calories	80
Fat	3.9 g
Carbohydrates	11 g
Protein	2 g
Dietary Fiber	2.5 g
Calcium	43 mg
Phosphorus	66 mg
Sodium	45 mg
Potassium	389 mg





LOW  
SODIUM



HIGH  
ACID

# Noodles Romano

Servings: 6

Serving size: 1/6 of recipe

## NUTRITION INFORMATION

(per serving)

**Calories** 380  
**Fat** 23.9 g  
**Carbohydrates** 32 g  
**Protein** 13 g  
**Dietary Fiber** 3.2 g  
**Calcium** 211 mg  
**Phosphorus** 237 mg  
**Sodium** 332 mg  
**Potassium** 157 mg



**Dialysis-  
friendly**  
option:

Individuals on dialysis should reduce cheese by half to reduce the phosphorus content of this recipe. Also, remember to take your phosphorus binders if they have been prescribed to you.

## INGREDIENTS

1/2 cup unsalted butter (divided)	1/8 teaspoon pepper
2 tablespoons dried parsley	8 ounces whole wheat spaghetti
1 teaspoon dried basil	1 garlic clove
6 ounces light or reduced-fat cream cheese	3/4 cup Parmesan cheese

## COOKING INSTRUCTIONS

1. Combine 1/4 cup butter, parsley flakes and basil.
2. Blend cream cheese and pepper in with butter mixture.
3. Stir in 2/3 cup boiling water. Blend mixture well.
4. Keep warm over pan of hot water.
5. Cook noodles in unsalted water until just tender; drain.
6. Cook minced garlic in 1/4 cup butter for 1 to 2 minutes, then pour over noodles, tossing lightly and quickly to coat well.
7. Sprinkle with 1/2 cup Parmesan cheese; toss again.
8. Pile noodles on warm serving platter.
9. Spoon warm cream cheese sauce over noodles.
10. Sprinkle with remaining 1/4 cup Parmesan cheese.
11. Garnish with additional parsley.

## SERVING SUGGESTION

A dish worthy of an entree, but make sure to serve with some type of vegetables on the side to balance out the acids from the cheese.





# Quinoa With Black Beans and Avocado

Servings: 6

Serving size: 1/6 of recipe



LOW SODIUM



LOW PROTEIN



REDUCES ACID

## INGREDIENTS

1 tablespoon olive oil  
 1 cup dry quinoa, rinsed  
 1 3/4 cups water  
 1 can (16 ounces) low-sodium black beans, drained and rinsed (rinse twice to reduce sodium)  
 1 avocado, chopped into chunks

3/4 cup cherry tomatoes, quartered  
 1/2 medium red onion, diced  
 1 small garlic clove, minced  
 1 medium red bell pepper, chopped into chunks  
 1/4 cup fresh cilantro, chopped

## Dressing:

1 lime, juiced  
 1/2 tablespoon olive oil  
 1/2 teaspoon cumin

## NUTRITION INFORMATION

(per serving)

Calories 260  
 Fat 9.1 g  
 Carbohydrates 37 g  
 Protein 9 g  
 Dietary Fiber 9.7 g  
 Calcium 62 mg  
 Phosphorus 230 mg  
 Sodium 8 mg  
 Potassium 596 mg

## COOKING INSTRUCTIONS

1. Warm the olive oil in a medium saucepan over medium heat. Add rinsed quinoa and toast for 2 to 3 minutes until it starts to smell nutty.
2. Add water, stir once, cover and simmer with a lid for 20 minutes.
3. While quinoa is cooking, prepare dressing by combining the lime juice, oil and cumin. Whisk aggressively. Adjust seasoning to taste.
4. When the quinoa is finished, remove from heat and fluff with a fork. Add black beans and toss to warm through.
5. Let the quinoa cool for about 5 minutes and then add all of the remaining ingredients, including dressing, and mix. Adjust seasoning if necessary.



**Dialysis-friendly**  
option:

If on dialysis and watching potassium intake, leave avocado out of recipe.





LOW SODIUM



LOW PROTEIN



REDUCES ACID

**NUTRITION INFORMATION**

(per serving)

Calories 95

Fat 3.6 g

Carbohydrates 16 g

Protein 2 g

Dietary Fiber 2.6 g

Calcium 29 mg

Phosphorus 47 mg

Sodium 46 mg

Potassium 381 mg

# Roasted Vegetables

Servings: 8

Serving size: 1/8 of recipe

Recipe submitted by Nicole Harr

## INGREDIENTS

- |   |                                       |
|---|---------------------------------------|
| 1 small butternut squash, cubed             | 1 tablespoon fresh thyme, chopped     |
| 2 medium red bell peppers, seeded and diced | 2 tablespoons fresh rosemary, chopped |
| 1 small sweet potato, peeled and cubed      | 3 tablespoons olive oil               |
| 2 small Yukon Gold potatoes, cubed          | 1½ tablespoons balsamic vinegar       |
| 1 medium red onion, quartered               | Salt to taste                         |
|   | Freshly ground black pepper           |

## COOKING INSTRUCTIONS

1. Preheat oven to 475° F.
2. In a large bowl, combine the squash, red bell peppers, sweet potato and Yukon Gold potatoes. Separate red onion quarters into pieces and add them to the mixture.
3. In a small bowl, stir together thyme, rosemary, olive oil, vinegar, salt and pepper.
4. Toss oil mixture with vegetables until they are coated.
5. Spread evenly on a large pan.
6. Roast for 35 to 40 minutes in the oven, stirring every 10 minutes or until vegetables are cooked through and browned.



# Couscous and Feta-Stuffed Peppers

Servings: 4

Serving size: 1 stuffed pepper



LOW  
SODIUM



HIGH  
FLUID



REDUCES  
ACID

## INGREDIENTS

Vegetable oil cooking spray	6 ounces yellow squash,
1¼ cups low-sodium chicken	quartered lengthwise, then
or vegetable broth	sliced across thinly
⅔ cup couscous	½ teaspoon fennel seeds
4 large bell peppers, mixed colors	½ teaspoon dried oregano
1¼ tablespoon olive oil	1 cup cherry tomatoes, cut in half
½ cup onion, chopped	15 ounces canned low-sodium
6 ounces zucchini, quartered	garbanzo beans, drained and
lengthwise, then sliced	rinsed
across thinly	4 ounces low-fat or reduced-fat
	crumbled feta cheese

## NUTRITION INFORMATION

(per serving)

Calories 400  
 Fat 12.1 g  
 Carbohydrates 57 g  
 Protein 19 g  
 Dietary Fiber 9.5 g  
 Calcium 151 mg  
 Phosphorus 272 mg  
 Sodium 385 mg  
 Potassium 797 mg

## COOKING INSTRUCTIONS

1. Preheat oven to 350° F.
2. Coat a small baking dish with cooking spray.
3. Bring the broth to a boil in a saucepan, add the couscous, cover the pan and remove it from the heat.
4. Meanwhile, boil water in a large pot. Cut the stems and top ½-inch off the bell peppers; scoop out the seeds and membranes. Boil trimmed peppers for 5 minutes, then drain them upside down.
5. Heat oil in a nonstick skillet. Add onion, zucchini, yellow squash, fennel seeds and oregano.
6. Cook, stirring frequently, for 5 minutes or until softened.
7. Remove from heat and stir in the tomatoes and garbanzo beans.
8. Using a fork, scrape the couscous into the skillet and toss with the vegetables.
9. Stir in the crumbled feta cheese.
10. Place peppers upright in the baking dish and fill them with couscous.
11. Bake 15 minutes. Serve immediately.



**Dialysis-  
friendly**  
option:

For individuals on dialysis and watching potassium and phosphorus intake, this recipe should be avoided.



# Entrees

All entrees with meat should be paired with vegetables to help balance out the dietary acids from the meat.

Grilled Salmon With Papaya-Mint Salsa	75
Rosemary Sage Burgers With Apple Slaw and Chive Mayo	76
Peach Pork Tenderloin	77
Salmon With Pineapple Salsa	78
Sweet Potato Burgers	79
Baked Salmon	80
Shrimp and Crab Gumbo	81
Chicken and Spanish Rice	83
Grilled Malt Chicken With Peppers and Onion	84
Summer Shrimp	85
Coffee-Ancho Chile-Rubbed Flat Iron Steak	86
Mushroom and Broccoli Risotto	87

Grilled Salmon  
With Papaya-Mint Salsa



# Grilled Salmon With Papaya-Mint Salsa

Servings: 4

Serving size: 1 salmon steak



LOW  
SODIUM



HIGH  
ACID

## INGREDIENTS

¼ cup papaya, peeled and chopped	1 tablespoon fresh lime juice
¼ cup yellow bell pepper, chopped	1 teaspoon fresh ginger, grated
¼ cup green onion, thinly sliced	1 teaspoon jalapeño pepper, seeded and minced
1 tablespoon pimiento, chopped	Vegetable oil cooking spray
1 tablespoon fresh mint, chopped	4 salmon steaks (4 ounces each) or fillets, about 1 to 1¼ inches thick
1 tablespoon rice wine or white vinegar	

## NUTRITION INFORMATION

(per serving)

**Calories** 195  
**Fat** 8.9 g  
**Carbohydrates** 3 g  
**Protein** 25 g  
**Dietary Fiber** 0.5 g  
**Calcium** 21 mg  
**Phosphorus** 345 mg  
**Sodium** 56 mg  
**Potassium** 533 mg

## COOKING INSTRUCTIONS

1. For salsa, combine all ingredients except the salmon in a small bowl.
2. Cover and chill at least 30 minutes.
3. Lightly coat grill or broiler pan with cooking spray.
4. Sprinkle both sides of salmon with pepper.
5. Grill or broil for 5 minutes on each side or until done.
6. Top each salmon steak with ¼ cup salsa.

## SERVING SUGGESTION

Pair with whole-wheat couscous and side salad or fruit dish.



LOW  
SODIUMHIGH  
ACID

# Rosemary Sage Burger With Apple Slaw and Chive Mayo

Servings: 4

Serving size: 1 burger

## NUTRITION INFORMATION

(per serving)

**Calories** 285**Fat** 12.5 g**Carbohydrates** 25 g**Protein** 19 g**Dietary Fiber** 2.6 g**Calcium** 77 mg**Phosphorus** 192 mg**Sodium** 356 mg**Potassium** 420 mg

## INGREDIENTS

### Burgers:

6 ounces 90 percent extra-lean ground round beef	1 teaspoon ground sage
6 ounces lean ground pork	¼ teaspoon salt
1 tablespoon fresh rosemary, chopped	4 whole-grain buns
	1 cup baby spinach leaves

### Chive mayo:

½ cup nonfat sour cream	1 dash salt
2 tablespoons chives, chopped	1 dash pepper

### Apple slaw:

3 medium green apples, peeled, cored and grated	2 teaspoons extra-virgin olive oil
	1 teaspoon fresh lemon juice

## COOKING INSTRUCTIONS

1. Preheat grill to medium-high.
2. Combine beef, pork, rosemary, sage and salt in a medium-size bowl; form into 4 patties.
3. In a small bowl, combine chive mayo ingredients.
4. In a medium bowl, combine slaw ingredients.
5. Grill patties 4 minutes on each side or until juices run clear.
6. Toast the buns and spread each top with the chive mayo. Place burger on bun, then top with slaw and several spinach leaves.

## SERVING SUGGESTION

Baked fries will help balance out the acids in the burger.



# Peach Pork Tenderloin

Servings: 8

Serving size: 1/8 of recipe



LOW  
SODIUM



HIGH  
ACID

## INGREDIENTS

2 pork tenderloins (1 pound each), trimmed	1/2 cup packed light brown sugar
Salt (optional)	1/2 cup water
Fresh ground black pepper	1 lemon, juiced
1/2 teaspoon olive oil	2 teaspoons whole-grain mustard
3 medium peaches (firm but ripe), pitted and wedged	1/4 teaspoon pepper
1/2 cup dry white wine or white wine vinegar	1/4 teaspoon fresh rosemary, chopped plus 1 to 2 additional springs

## COOKING INSTRUCTIONS

1. Preheat grill to high.
2. Pat pork dry and season with pepper (and salt lightly, if you wish).
3. Grill pork on lightly oiled grill until marked, about 10 minutes on each side.
4. While the meat is cooking, combine peaches, wine/vinegar, brown sugar and water in a saucepan.
5. Peel strips of lemon skin into pan and squeeze in the juice.
6. Cover and cook on medium to high heat until peaches are tender, about 5 minutes.
7. Move solid ingredients to another bowl with a slotted spoon.
8. Add mustard, pinch of salt and 1/4 teaspoon pepper to the juice in the pan.
9. Move 1/3 cup of the juice to a bowl, add chopped rosemary and brush onto pork; grill pork for additional 10 to 15 minutes, until thermometer in the thickest part of the meat shows 145° F. Transfer pork to cutting board to cool.
10. Simmer remaining liquid in saucepan for a few minutes until it thickens; remove from heat, add peaches and rosemary sprigs.
11. Slice pork and serve with peach sauce.

### NUTRITION INFORMATION

(per serving)

Calories	190
Fat	4.4 g
Carbohydrates	15 g
Protein	22 g
Dietary Fiber	0.9 g
Calcium	19 mg
Phosphorus	277 mg
Sodium	85 mg
Potassium	580 mg



**Dialysis-friendly**  
option:

For individuals on dialysis watching potassium intake, this recipe should be avoided.



LOW  
SODIUMHIGH  
ACID

# Salmon With Pineapple Salsa

Servings: 6

Serving size: 4 ounces

## NUTRITION INFORMATION

(per serving)

**Calories** 185**Fat** 7.2 g**Carbohydrates** 8 g**Protein** 22 g**Dietary Fiber** 0.4 g**Calcium** 19 mg**Phosphorus** 297 mg**Sodium** 180 mg**Potassium** 457 mg

## INGREDIENTS

1½ pounds salmon fillets

### Marinade:

2 tablespoons low-sodium

soy sauce

½ clove garlic, minced

¼ teaspoon ground ginger

2 tablespoons brown sugar

¼ cup unsweetened pineapple

juice

1½ tablespoons apple cider

vinegar

### Pineapple salsa:

¼ cup red bell pepper, finely  
chopped2 tablespoons fresh cilantro,  
finely chopped2 tablespoons red onion, finely  
chopped

1 cup fresh pineapple, chopped

## COOKING INSTRUCTIONS

1. In small bowl, make marinade by mixing soy sauce, garlic, ginger, brown sugar, pineapple juice and apple cider vinegar.
2. Measure out 2 tablespoons and set aside for pineapple salsa.
3. Place salmon in large resealable plastic bag or glass dish. Add the rest of the marinade minus 2 tablespoons (held aside); turn to coat well. Refrigerate 15 minutes or longer for extra flavor.
4. Make pineapple salsa by mixing 2 tablespoons of the marinade, pineapple, bell pepper, cilantro and onion in small bowl until well blended.
5. Remove salmon from marinade. Discard remaining marinade.
6. Grill salmon over medium-high heat 6 to 7 minutes per side or until fish flakes easily with a fork, brushing with remaining 2 tablespoons marinade halfway through cooking.
7. Serve salmon with pineapple salsa.





# Sweet Potato Burgers

Servings: 4

Serving size: 1 burger

Recipe submitted by Hannah Harnden, 23 years old with PKD, first generation



## INGREDIENTS

1 medium sweet potato, peeled and cubed	Pinch of salt
1 can (15.5 ounces) low-sodium black beans	1/3 cup all-purpose flour
1/2 cup onion, chopped	1 cup greens
3/4 teaspoon apple cider vinegar	1 cup panko for coating
1 teaspoon garlic powder	1/4 cup olive oil
1/2 teaspoon chili powder	4 whole-grain buns
	Toppings: Avocado, spicy mustard, greens and tomato

## COOKING INSTRUCTIONS

1. Cook potato and mash together with black beans.
2. Add the onion, vinegar, garlic powder, chili powder, salt, flour and greens; continue to mash with a fork.
3. Form into 4 patties and coat with panko.
4. Fry in pan and serve with toppings of your choice.

### NUTRITION INFORMATION

(per serving)

Calories	365
Fat	6.8 g
Carbohydrates	62 g
Protein	15 g
Dietary Fiber	12.9 g
Calcium	126 mg
Phosphorus	247 mg
Sodium	286 mg
Potassium	632 mg



**Dialysis-friendly option:**

If on dialysis and watching potassium intake, consider first boiling the potato based on instructions on page 19 and do not top with avocado.



LOW  
SODIUMHIGH  
ACID

# Baked Salmon

Servings: 4      Serving size: 1 fillet

Recipe submitted by Nicole Harr

## INGREDIENTS

2 cloves garlic, minced	½ tablespoon fresh lemon juice
3 tablespoons light olive oil	½ tablespoon fresh parsley, chopped
1 teaspoon dried basil	4 4-ounce salmon fillets
Pinch of salt (optional)	
¼ teaspoon ground black pepper	

## NUTRITION INFORMATION

(per serving)

**Calories** 245

**Fat** 17.3 g

**Carbohydrates** 1 g

**Protein** 21 g

**Dietary Fiber** 0.2 g

**Calcium** 19 mg

**Phosphorus** 291 mg

**Sodium** 83 mg

**Potassium** 416 mg

## COOKING INSTRUCTIONS

1. In a medium glass bowl, prepare marinade by mixing garlic, light olive oil, basil, salt, pepper, lemon juice and parsley.
2. Place salmon fillets in a medium glass baking dish and cover with the marinade.
3. Marinate in the refrigerator about 1 hour, turning occasionally.
4. Preheat oven to 375° F.
5. Place fillets in aluminum foil, cover with marinade and seal.
6. Place sealed salmon in the glass dish and bake 35 to 45 minutes, until easily flaked with a fork.



# Shrimp and Crab Gumbo

Servings: 6

Serving size: 2 cups



LOW  
SODIUM



HIGH  
FLUID



HIGH  
ACID

## INGREDIENTS

¼ cup olive oil	8 ounces uncooked fresh shrimp
⅓ cup all-purpose flour	6 ounces crab meat (can also use imitation crab meat)
1 cup bell pepper, chopped	1 cup green onion tops, diced
1½ cups onion, diced	¼ cup fresh parsley, chopped
1 clove garlic, chopped	Pepper to taste
1 tablespoon celery leaves or diced celery	1 teaspoon hot sauce (optional)
4 cups water (divided use)	6 cups cooked white rice (1 cup mixed in; 5 cups for serving gumbo over)
4 cups low-sodium chicken broth (see page 108)	

## NUTRITION INFORMATION

(per serving)

Calories	415
Fat	11.2 g
Carbohydrates	60 g
Protein	18 g
Dietary Fiber	2.0 g
Calcium	75 mg
Phosphorus	283 mg
Sodium	501 mg
Potassium	418 mg

## COOKING INSTRUCTIONS

1. Make a roux by heating oil and flour in a large skillet over medium heat. Stir constantly until flour is pecan-colored.
2. Add bell pepper, onion, garlic, celery leaves and 1 cup water.
3. Cover and cook on very low heat until vegetables are tender.
4. Increase heat to high, and add 3 cups water and 4 cups low-sodium chicken broth. Boil for 5 minutes.
5. Reduce to medium heat, and add shrimp and crab meat. Boil 10 minutes.
6. Add green onion tops and parsley.
7. Reduce to low heat, and simmer for 5 minutes.
8. Add 1 cup cooked rice.
9. Season with pepper and hot sauce.
10. Serve gumbo over hot rice.

## NOTES

Frozen shrimp are often high in sodium.





# Chicken and Spanish Rice

Servings: 5

Serving size: 1/5 of recipe



LOW  
SODIUM



HIGH  
FLUID



HIGH  
ACID

## INGREDIENTS

1 cup onions, chopped	1/2 teaspoon black pepper
3/4 cup green pepper, chopped	1/4 teaspoon garlic, minced
1 tablespoon olive oil	5 cups cooked brown rice (cooked in unsalted water)
1 can (8 ounces) low-sodium tomato sauce	2 1/2 cups chicken breasts, cooked, skin and bone removed, diced
1 teaspoon fresh parsley, chopped	

## COOKING INSTRUCTIONS

1. In a large skillet, sauté onions and green peppers in oil for 5 minutes on medium heat.
2. Add tomato sauce and spices. Heat through.
3. Add cooked rice and chicken. Heat through.

### NUTRITION INFORMATION

(per serving)

**Calories** 360

**Fat** 7.0 g

**Carbohydrates** 51 g

**Protein** 23 g

**Dietary Fiber** 6.7 g

**Calcium** 51 mg

**Phosphorus** 300 mg

**Sodium** 57 mg

**Potassium** 379 mg



LOW  
SODIUMHIGH  
ACID

# Grilled Malt Chicken With Peppers and Onion

*Servings: 8**Serving size: 3 ounces**Recipe submitted by Geno Leage, PKD chapter volunteer*

## NUTRITION INFORMATION

(per serving)

**Calories** 155**Fat** 6.3 g**Carbohydrates** 4 g**Protein** 20 g**Dietary Fiber** 0.9 g**Calcium** 23 mg**Phosphorus** 156 mg**Sodium** 215 mg**Potassium** 276 mg

## INGREDIENTS

- |  |   |
|--|---|
| 4 4-ounce chicken breasts,<br>skinless | 4 ounces mushrooms, Crimini<br>or button, washed and sliced |
| 1 medium yellow bell pepper,<br>sliced | 3 cloves garlic, minced                                     |
| 1 medium red bell pepper, sliced       | ½ teaspoon dried oregano                                    |
| 1 medium onion, sliced                 | Pinch salt and pepper, to taste                             |

### Marinade:

- |                            |                                   |
|----------------------------|-----------------------------------|
| 3 tablespoons malt vinegar | ½ teaspoon salt                   |
| ¼ cup olive oil            | ½ teaspoon black ground<br>pepper |
| 1 shallot sliced           |                                   |
| 1 teaspoon dried oregano   |                                   |

## COOKING INSTRUCTIONS

1. Heat grill to medium-high heat.
2. Combine all marinade ingredients and mix well. Place the chicken and marinade in a resealable bag. Marinate the chicken for 30 to 45 minutes.
3. Remove chicken and let drain for a moment. Place on the grill to cook.
4. Sauté the onion and pepper in a separate pan. Cook for about 5 minutes, then add mushrooms. Cook until all vegetables are soft.
5. Turn chicken on the grill, cook until cooked thoroughly.
6. Place the chicken on the plate first and cover with the vegetables.



# Summer Shrimp

Servings: 4

Serving size: 2 skewers



LOW  
SODIUM



LOW  
PROTEIN



HIGH  
ACID

## INGREDIENTS

### Marinade:

2 cloves garlic, smashed	1 teaspoon low-sodium
1 tablespoon olive oil	Worcestershire sauce
2 teaspoons fresh lime juice	¼ teaspoon freshly ground
1 teaspoon dried oregano	black pepper

### Shrimp:

1 small red onion	24 large fresh shrimp (about
8 bamboo skewers, 10 inches long,	1 pound), peeled and deveined
soaked in water for 30 minutes	16 serrano chilies

### NUTRITION INFORMATION

(per serving)

Calories 85

Fat 3.6 g

Carbohydrates 4 g

Protein 9 g

Dietary Fiber 1.3 g

Calcium 40 mg

Phosphorus 104 mg

Sodium 48 mg

Potassium 218 mg

## COOKING INSTRUCTIONS

1. In a blender or mini food processor, process all marinade ingredients until smooth.
2. Toss shrimp with marinade in a bowl to coat well. Set aside at least 10 minutes or up to 2 hours.
3. Cut onion in half and each half into quarters.
4. Thread 1 chili onto each skewer. Add 1 shrimp and a chunk of onion; repeat twice. Finish each skewer with a second chili.
5. Heat a cast-iron grill pan or an electric or gas grill over high heat.
6. Cook skewers, covered, turning once, until shrimp are cooked through and onions are charred and just tender, about 5 minutes per side.
7. Serve with salsa.

## NOTES

If using metal skewers, spray with a vegetable oil cooking spray. If using a charcoal grill, be sure to adjust cooking time because the shrimp will cook faster. Frozen shrimp are often high in sodium.





LOW  
SODIUM



HIGH  
ACID

# Coffee-Ancho Chile- Rubbed **Flat Iron Steak**

Servings: 4

Serving size: 1 steak

Recipe submitted by Geno Leage, PKD chapter volunteer

## INGREDIENTS

4 4-ounce flat iron steaks	½ cup chicken stock
Coffee-Ancho Chile Rub (see recipe on page 105)	1 teaspoon cornstarch
	1½ teaspoons water

### NUTRITION INFORMATION

(per serving)

Calories 290

Fat 21.1 g

Carbohydrates 5 g

Protein 20 g

Dietary Fiber 0.9 g

Calcium 17 mg

Phosphorus 169 mg

Sodium 155 mg

Potassium 365 mg

Caffeine 7 mg

## COOKING INSTRUCTIONS

1. Generously rub steaks with spice rub and let sit for 30 minutes before cooking.
2. Cook to your preferred doneness.
3. Remove from pan, but leave the pan on the burner and add the stock.
4. Combine cornstarch and water.
5. Bring the stock to a quick boil, add the cornstarch slurry and whisk. The stock will thicken quickly; remove from heat and serve over the steak.

## SERVING SUGGESTION

Serve with potato and vegetable side dishes from the *Sides* section starting on page 62.





# Mushroom and Broccoli Risotto

Servings: 4

Serving size: ¼ of recipe

Recipe submitted by Geno Leage, PKD chapter volunteer



LOW  
SODIUM



LOW  
PROTEIN



HIGH  
FLUID

## INGREDIENTS

2 - 2½ cups chicken stock (see recipe on page 108)	1 tablespoon olive oil
8 ounces button mushrooms, washed and sliced	1 cup Arborio rice
½ medium onion, diced	⅓ cup white wine
2 cloves garlic, minced	2 tablespoons Parmesan cheese, grated
¼ pound broccoli florets, cut into bite-size pieces	Pepper to taste

## COOKING INSTRUCTIONS

1. Heat chicken stock.
2. Prep vegetables and measure out dry goods. Once the stock comes to a boil, remove from heat and set aside.
3. Heat olive oil in medium-size saucepan; add onion, mushrooms and garlic. Sauté until onions become soft.
4. Add the risotto and the rice and cook for a few minutes.
5. Add the wine and reduce by half.
6. Add 1 cup of the hot stock; stir rice continuously as it absorbs the stock. Continue adding more stock and stirring.
7. After all stock is added, mix in the broccoli.
8. After the stock is absorbed and the rice is tender, add Parmesan cheese and season to taste.

### NUTRITION INFORMATION

(per serving)

Calories 300

Fat 5.8 g

Carbohydrates 49 g

Protein 10 g

Dietary Fiber 2.5 g

Calcium 76 mg

Phosphorus 189 mg

Sodium 136 mg

Potassium 419 mg



# Desserts



Strawberry-Mango Parfaits With Ginger Topping	89
Chocolate Chip Cookies	90
Strawberry Lime Dessert	91
Light and Creamy Cranberry Fresh Fruit Dip	92
Blackberry Cobbler	93
Coconut Angel Food Cake	94
Lemon Ricotta Pie	96
Bread Pudding	97
Bran Muffin	98
Apple Muffin	99
Key Lime Pie	100
Zesty Lemon Tart	101

**Strawberry-Mango Parfaits  
With Ginger Topping**



# Strawberry-Mango Parfaits

## With Ginger Topping

Servings: 4

Serving size: 1 parfait



LOW  
SODIUM



LOW  
PROTEIN



HIGH  
FLUID



REDUCES  
ACID

### INGREDIENTS

1 pint fresh strawberries, stemmed and sliced	1 pint light or reduced-fat strawberry ice cream
2 tablespoons granulated sugar or to taste	½ cup gingersnap crumbs
2 tablespoons orange juice	2 tablespoons crystallized ginger, coarsely chopped (optional)
2 teaspoons orange rind, grated	
1 pint mango or orange sorbet	

### COOKING INSTRUCTIONS

1. Toss strawberries, sugar, juice and rind in a bowl; set aside.
2. Place scoops of sorbet and ice cream in 4 deep bowls or parfait glasses.
3. Top with a layer of sauce; repeat with another layer of sorbet and ice cream.
4. Top with remaining sauce, crumbs and ginger.

#### NUTRITION INFORMATION

(per serving)

**Calories** 380

**Fat** 7.2 g

**Carbohydrates** 78 g

**Protein** 4 g

**Dietary Fiber** 2.9 g

**Calcium** 128 mg

**Phosphorus** 106 mg

**Sodium** 142 mg

**Potassium** 351 mg



## DESSERTS



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

### NUTRITION INFORMATION

(per serving)

Calories 230

Fat 19.1 g

Carbohydrates 13 g

Protein 5 g

Dietary Fiber 2.8 g

Calcium 52 mg

Phosphorus 110 mg

Sodium 128 mg

Potassium 192 mg

Caffeine 6 mg

# Chocolate Chip Cookies

Servings: 6

Serving size: 1 cookie

Recipe submitted by Emily Decker

## INGREDIENTS

1 cup blanched almond flour, sifted	2 tablespoons honey
Sprinkle of sea salt ( $\frac{1}{4}$ teaspoon or less to taste)	1 $\frac{1}{2}$ teaspoons vanilla
$\frac{1}{8}$ teaspoon baking soda	$\frac{1}{4}$ cup (or more for extra chocolatey goodness) Enjoy Life Chocolate Chips or chopped dark chocolate (70-85 percent or higher)
$\frac{1}{4}$ teaspoon cinnamon	
3 tablespoons melted coconut oil	

## COOKING INSTRUCTIONS

1. Preheat oven to 350° F.
2. Combine almond flour, salt, baking soda and cinnamon in a large mixing bowl or stand mixer; mix thoroughly.
3. Combine liquid ingredients in a medium mixing bowl.
4. Slowly add liquid ingredients to the dry ingredients, mixing well until combined.
5. Stir in chocolate chips.
6. Spoon onto prepared baking sheet and bake for 10 to 11 minutes or until golden brown around the edges.



# Strawberry Lime Dessert

Servings: 6

Serving size: 1 glass or dish



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

## INGREDIENTS

4 cups fresh strawberries, remove stems and quarter	1 tablespoon sugar
½ cup whipping cream	¼ teaspoon lime peel zest
	1 lime, cut into 6 to 8 wedges

## COOKING INSTRUCTIONS

1. In a shallow dish, mash 1 cup of the quartered strawberries with a potato masher, or puree with a mini food chopper, food processor or blender.
2. In a medium bowl, beat whipping cream, sugar and lime peel until cream is very stiff.
3. Fold in mashed strawberries.
4. In 6 individual 6- to 7-ounce glasses or dishes, layer whipped cream mixture and remaining strawberries.
5. Top with a fresh lime wedge. Serve immediately or cover and chill up to 2 hours before serving.

### NUTRITION INFORMATION

(per serving)

Calories 90

Fat 6.4 g

Carbohydrates 9 g

Protein 1 g

Dietary Fiber 1.5 g

Calcium 26 mg

Phosphorus 30 mg

Sodium 8 mg

Potassium 136 mg



## DESSERTS



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

# Light and Creamy Cranberry **Fresh Fruit Dip**

Servings: 12

Serving size: 2 tablespoons

### NUTRITION INFORMATION

(per serving)

Calories 45

Fat 1.2 g

Carbohydrates 7 g

Protein 1 g

Dietary Fiber 0.1 g

Calcium 25 mg

Phosphorus 25 mg

Sodium 21 mg

Potassium 47 mg

### INGREDIENTS

8 ounces light sour cream

¼ teaspoon nutmeg

½ cup whole-berry cranberry

¼ teaspoon ground ginger

sauce

### COOKING INSTRUCTIONS

1. Put all ingredients in a blender or food processor; blend until well mixed.
2. Serve with your favorite fresh fruit such as pineapple, pears and strawberries.



# Blackberry Cobbler

Servings: 6

Serving size: 1/6 of recipe



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

## INGREDIENTS

Vegetable oil cooking spray	2/3 cup firmly packed brown sugar
1 12-ounce bag frozen unsweetened blackberries, thawed (or 3 half-pint containers of fresh blackberries)	3/4 cups quick oats
1/4 cup granulated sugar	1/2 cup all-purpose flour
1 1/2 tablespoons cornstarch	1/2 teaspoon ground cinnamon
1 1/2 tablespoons water	1/2 cup unsalted butter, softened
	1 8-ounce container light or reduced-fat whipped cream

## COOKING INSTRUCTIONS

1. Heat oven to 350° F.
2. Lightly grease an 8-inch square baking dish.
3. In large saucepan, combine blackberries and sugar, let stand for 30 minutes.
4. Cook blackberries and sugar mixture, stirring frequently, over medium heat for 5 minutes.
5. In a small bowl, combine cornstarch and water until blended.
6. Stir cornstarch mixture into blackberry mixture.
7. Bring to boil over medium heat; boil for 1 minute or until mixture is thickened.
8. Spoon mixture into pan.
9. In a medium bowl, combine brown sugar, oats, flour and cinnamon.
10. Using a pastry blender or fork, cut in butter until mixture is crumbly. Sprinkle evenly over blackberry mixture.
11. Bake 30 to 40 minutes or until lightly browned and bubbly.
12. Serve with a dollop of whipped cream.

### NUTRITION INFORMATION

(per serving)

Calories 430

Fat 18.2 g

Carbohydrates 65 g

Protein 4 g

Dietary Fiber 3.4 g

Calcium 55 mg

Phosphorus 77 mg

Sodium 21 mg

Potassium 156 mg





LOW SODIUM



LOW PROTEIN



REDUCES ACID

# Coconut Angel Food Cake

Servings: 12

Serving size: 1 slice and 3 tablespoons sauce

## INGREDIENTS

### Cake:

¼ teaspoon salt  
 1 teaspoon cream of tartar  
 1 cup superfine sugar  
 2 teaspoons pure vanilla extract  
 2 teaspoons fresh lemon juice  
 1 cup sifted cake flour

⅓ cup dry, unsweetened, shredded coconut  
 ⅓ cup fresh coconut shavings  
 1¼ cups egg whites (from about 9 large eggs), at room temperature

### Sauce:

1 bag (12 ounces) cranberries, fresh or frozen and thawed  
 ⅓ cup granulated sugar  
 1 tablespoon grated orange zest

⅔ cup fresh orange juice  
 2 tablespoons orange-flavored liqueur  
 ½ teaspoon cinnamon

### NUTRITION INFORMATION

(per serving)

Calories 215  
 Fat 2.4 g  
 Carbohydrates 44 g  
 Protein 4 g  
 Dietary Fiber 2.2 g  
 Calcium 10 mg  
 Phosphorus 27 mg  
 Sodium 94 mg  
 Potassium 167 mg





## **COOKING INSTRUCTIONS**

### **Cake:**

1. Heat oven to 325° F.
2. Beat egg whites and salt in a bowl with an electric mixer on medium until frothy, about 2 minutes.
3. Add cream of tartar. Beat until soft peaks form, about 5 minutes.
4. Increase speed to high. Add ½ cup sugar, 1 tablespoon at a time, until whites form soft, glossy peaks.
5. Beat in vanilla and lemon juice.
6. In another bowl, mix remaining ½ cup sugar with flour.
7. Sift ⅓ sugar/flour mixture over egg whites; gently, but thoroughly, fold into batter using a large spatula. Repeat twice with remaining sugar/flour mixture.
8. Fold in dry coconut in 2 batches.
9. Scrape batter into an ungreased 10-inch angel food tube pan with a removable bottom.
10. Smooth top into even layer.
11. Bake 35 minutes or until top is lightly browned and cake springs back when you press it.
12. Turn pan upside down and rest on the neck of a glass bottle to cool.
13. Run a knife against insides of pan to loosen cake; remove pan. Loosen cake from bottom of pan; invert onto a plate.
14. Cover with plastic wrap. Store at room temperature or in the refrigerator.

### **Sauce:**

1. Boil all ingredients in a medium saucepan 1 minute, until lightly thickened.
2. Let cool.
3. Cover; refrigerate 1 hour.
4. Before serving, sprinkle top of cake with coconut shavings and cut into 12 slices. Serve slices with 3 tablespoons sauce.



## DESSERTS



LOW  
SODIUM



LOW  
PROTEIN

# Lemon Ricotta Pie

Servings: 8

Serving size: 1 slice

### NUTRITION INFORMATION

(per serving)

Calories 105

Fat 6.2 g

Carbohydrates 4 g

Protein 8 g

Dietary Fiber 0.1 g

Calcium 156 mg

Phosphorus 130 mg

Sodium 127 mg

Potassium 98 mg

### INGREDIENTS

3 large eggs, separated  
15 ounces ricotta cheese  
(part-skim; whole can be  
used as well)  
Juice and zest of 1 medium  
lemon

$\frac{2}{3}$  cup sugar substitute  
(Splenda)  
 $\frac{1}{2}$  teaspoon vanilla  
Dash of salt

### COOKING INSTRUCTIONS

1. Preheat oven to 350° F.
2. Beat egg whites until soft peaks form.
3. Combine egg yolks, ricotta, lemon juice and zest, sugar substitute, vanilla and salt; stir to mix.
4. Test to make sure there is enough lemon flavor. If not, add more.
5. Fold egg whites into ricotta mixture and pour into greased pie pan.
6. Bake 25 minutes or until a toothpick in the middle comes out clean.
7. Refrigerate until completely cool.
8. Cut into 8 piece and serve with fresh berries or your favorite topping.

### SERVING SUGGESTION

Serving with berries, or other fruit, makes this a great dessert!



# Bread Pudding

Servings: 8

Serving size: 1/8 of recipe



LOW  
SODIUM



LOW  
PROTEIN

## INGREDIENTS

Vegetable oil cooking spray	1/2 cup liquid egg substitute
8 slices of bread or 2 large bagels	1/2 cup sugar
1 cup liquid nondairy low-fat or reduced-fat creamer	1 teaspoon cinnamon

## COOKING INSTRUCTIONS

1. Preheat oven to 350° F.
2. Spray a 1½-quart baking dish with cooking spray.
3. Break bread into small pieces and put in dish.
4. Mix creamer, egg substitute, sugar and cinnamon together and pour over bread.
5. Gently press with a fork so all the liquid is absorbed by the bread.
6. Bake for 30 minutes until brown on top.

### NUTRITION INFORMATION

(per serving)

**Calories** 145

**Fat** 2.1 g

**Carbohydrates** 28 g

**Protein** 4 g

**Dietary Fiber** 1.0 g

**Calcium** 74 mg

**Phosphorus** 51 mg

**Sodium** 170 mg

**Potassium** 109 mg





# Bran Muffin

Servings: 12

Serving size: 1 muffin

## NUTRITION INFORMATION

(per serving)

Calories 120

Fat 1.9 g

Carbohydrates 27 g

Protein 5 g

Dietary Fiber 2.9 g

Calcium 96 mg

Phosphorus 175 mg

Sodium 100 mg

Potassium 159 mg

## INGREDIENTS

1 cup all-purpose flour

2 teaspoons baking powder

½ teaspoon cinnamon

2 cups 100 percent oat bran

1¼ cups low-fat milk

½ cup packed brown sugar

1 large egg

½ cup unsweetened applesauce

## COOKING INSTRUCTIONS

1. Preheat oven to 400° F.
2. Spray muffin pan with cooking spray.
3. Mix bran, milk and sugar in another bowl and let stand 5 minutes.
4. Mix flour, baking powder, baking soda and cinnamon in a large bowl.
5. In the bowl with the bran, stir in egg and applesauce.
6. Add to flour mixture and stir until just moistened. Batter will be lumpy.
7. Spoon batter into muffin pan, filling each cup ¾ full.
8. Bake for 18 to 20 minutes or until golden brown.



**Dialysis-friendly**  
option:

Individuals on dialysis should avoid this recipe and instead make Apple Muffins found on page 99 which is also suitable for those living with PKD.



# Apple Muffin

Servings: 12

Serving size: 1 muffin



LOW  
SODIUM



LOW  
PROTEIN

## INGREDIENTS

2 eggs	1 teaspoon baking soda
1 cup Splenda (sugar also can be used)	1½ teaspoon cinnamon
½ cup canola oil	1¾ cup raw apple, peeled and cut into small pieces (other low-potassium fruit also can be used)
¼ cup water	
1 tablespoon vanilla	
1½ cups all-purpose white flour	

## COOKING INSTRUCTIONS

1. Preheat oven to 400° F.
2. Grease muffin pan or use 12 muffin liners.
3. Beat eggs in a large bowl. Add Splenda or sugar, oil and water; mix well.
4. Add vanilla.
5. In a separate bowl, combine flour, baking soda and 1 teaspoon cinnamon.
6. Stir flour mixture into egg mixture. Batter will be lumpy.
7. Fold in apple pieces.
8. Fill muffin cups three quarters full.
9. Mix remaining ½ teaspoon cinnamon with 1 teaspoon Splenda or sugar. Sprinkle on top of muffins.
10. Bake for 20 minutes or until lightly browned.

## SERVING SUGGESTION

Pair with a glass of juice or fruit smoothie for breakfast, and you're off to a great start.

### NUTRITION INFORMATION

(per serving)

Calories	165
Fat	10.1 g
Carbohydrates	15 g
Protein	3 g
Dietary Fiber	0.9 g
Calcium	11 mg
Phosphorus	33 mg
Sodium	116 mg
Potassium	47 mg



LOW  
SODIUMLOW  
PROTEIN

# Key Lime Pie

Servings: 10

Serving size: 1 slice

## INGREDIENTS

### Crust:

1½ cups graham crackers,  
crushed

¼ cup sugar

5 tablespoons unsalted butter,  
melted

### Filling:

1 can (14 ounces) nonfat,  
sweetened, condensed milk

½ cup fresh lime juice

1 tablespoon lime zest, minced

1 large egg

2 large eggs, yolks and whites  
separated

¼ teaspoon cream of tartar

2 cups low-calorie whipped  
cream (such as Cool Whip)

1 lime, thinly sliced (optional)

### NUTRITION INFORMATION

(per serving)

**Calories** 260**Fat** 8.1 g**Carbohydrates** 42 g**Protein** 6 g**Dietary Fiber** 0.6 g**Calcium** 137 mg**Phosphorus** 151 mg**Sodium** 153 mg**Potassium** 228 mg

## COOKING INSTRUCTIONS

1. Preheat oven to 325° F.
2. For crust: Mix graham cracker crumbs with sugar and butter. Spread evenly over bottom and sides of a nonstick 9-inch pie pan, and bake 5 minutes. Remove from oven; let cool.
3. For filling: In a medium bowl, using a fork, beat milk, lime juice and zest. Add 1 whole egg and 2 egg yolks (reserve whites) and beat well. Set aside.
4. In a large mixing bowl, beat egg whites until foamy. After about 20 seconds, add cream of tartar. Fold whites into custard mixture and pour into partially baked crust.
5. Bake 45 to 60 minutes or until set. Let cool.
6. Cut into 10 slices and serve with whipped cream. Garnish with lime slices, if desired.



# Zesty Lemon Tart

Servings: 1

Serving size: 1 mini pie

Recipe submitted by Pat Smithson



## INGREDIENTS

Individual graham cracker pie crusts	2 tablespoons light whipped cream
2.8 ounces (one third of an 8-ounce carton) nonfat lemon yogurt	Lemon zest

## COOKING INSTRUCTIONS

1. Fill pie crusts with yogurt, then light whipped cream.
2. Garnish with lemon zest.

## SERVING SUGGESTIONS

You can substitute any yogurt. Try key lime yogurt with lime zest or strawberry yogurt with fresh strawberries on top.

### NUTRITION INFORMATION

(per serving)

Calories 195

Fat 7.0 g

Carbohydrates 29 g

Protein 5 g

Dietary Fiber 0.7 g

Calcium 154 mg

Phosphorus 138 mg

Sodium 159 mg

Potassium 211 mg



# Spices, Spreads and Rubs

Homemade Salt-Free Mexican Spice Mix	<b>103</b>
Homemade Mayonnaise	<b>104</b>
Coffee-Ancho Chile Rub	<b>105</b>
Simple Kebob Marinade	<b>106</b>
Roasted Garlic	<b>107</b>
Basic Chicken Stock	<b>108</b>

**Homemade Salt-Free  
Mexican Spice Mix**





# Homemade Salt-Free Mexican Spice Mix

Servings: 8

Serving size: 1 tablespoon



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

## INGREDIENTS

2 tablespoons chili powder                      5 teaspoons paprika  
4½ teaspoons cumin                                3 teaspoons onion powder  
2½ teaspoons garlic powder

## COOKING INSTRUCTIONS

1. Mix all ingredients together and store in an airtight container.
2. Add a few tablespoons to chicken, lean ground beef or to add a Mexican flair to any dish.

## NOTES

Try smoked paprika for a warm, smoky flavor.

### NUTRITION INFORMATION

(per serving)

Calories 20

Fat 0.7 g

Carbohydrates 4 g

Protein 1 g

Dietary Fiber 1.5 g

Calcium 24 mg

Phosphorus 23 mg

Sodium 37 mg

Potassium 112 mg





LOW  
SODIUM



LOW  
PROTEIN

# Homemade Mayonnaise

Servings: 24

Serving size: 1 tablespoon

## INGREDIENTS

- |   |  |
|---|--|
| 1 large pasteurized egg at room temperature | A pinch or two of sugar, if desired  |
| 1¼ cup olive oil, divided                   | 2 to 3 teaspoons fresh lemon juice (or white wine vinegar or a mix of the two) |
| ½ teaspoon dry mustard                      |  |
| ¼ teaspoon salt (or less, to taste)         |  |

### NUTRITION INFORMATION

(per serving)

Calories 105

Fat 11.5 g

Carbohydrates 0 g

Protein 0 g

Dietary Fiber 0 g

Calcium 1 mg

Phosphorus 4 mg

Sodium 27 mg

Potassium 4 mg

## COOKING INSTRUCTIONS

1. Put egg, ¼ cup oil, dry mustard, salt and sugar (if desired) in blender, food processor or mixer. Mix thoroughly.
2. With mixer running, very slowly (this is key) drizzle in remaining oil, a few drops at a time, allowing the oil to be completely absorbed by the mixture before adding more.
3. After all of the oil is added and the mixture has emulsified (it will seem to thicken and get lighter), add lemon juice/vinegar, stirring with spoon to incorporate.
4. Chill for 1 to 2 hours and then serve. Mayonnaise will keep for about a week in the refrigerator.

## NOTES

Because a raw egg is used, only use the freshest grade A or AA eggs with intact shells. Ideally, use shell eggs that have been pasteurized or otherwise treated to destroy salmonella.



# Coffee-Ancho Chile Rub

Servings: 14      Serving size: 1 tablespoon

Recipe submitted by Geno Leage, PKD chapter volunteer

## INGREDIENTS

¼ cup espresso coffee beans, finely ground	1 teaspoon dry mustard
⅓ cup packed brown sugar	2 teaspoons salt
1 tablespoon paprika	1 teaspoon black pepper
1 tablespoon garlic, granulated	1 tablespoon ancho chile powder
1 teaspoon dried oregano	

## COOKING INSTRUCTIONS

1. Grind coffee beans to a fine consistency.
2. Remove from grinder and place in a small mixing bowl.
3. Place the rest of the ingredients into the bowl and mix well.
4. This can be stored in a baggy or an airtight container.

## SERVING SUGGESTION

To be used with beef or lamb.



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

## NUTRITION INFORMATION

(per serving)

Calories 30  
 Fat 0.2 g  
 Carbohydrates 7 g  
 Protein 0 g  
 Dietary Fiber 0.6 g  
 Calcium 10 mg  
 Phosphorus 9 mg  
 Sodium 340 mg  
 Potassium 73 mg  
 Caffeine 27 mg





LOW  
SODIUM



LOW  
PROTEIN

# Simple Kebob Marinade

Servings: 8

Serving size: 2 tablespoons

## INGREDIENTS

¼ cup low-sodium soy sauce	1 clove crushed garlic
2 tablespoons fresh lemon juice	2 tablespoons honey or agave
	¾ cup olive oil

## NUTRITION INFORMATION

(per serving)

Calories 200  
Fat 20.3 g  
Carbohydrates 5 g  
Protein 1 g  
Dietary Fiber 0.1 g  
Calcium 3 mg  
Phosphorus 10 mg  
Sodium 266 mg  
Potassium 23 mg

## COOKING INSTRUCTIONS

1. Mix all ingredients together, adding oil last.
2. Marinate meat (chicken, pork, lean beef) or vegetables (onions, mushrooms, bell peppers, tomatoes, squash, cauliflower and more) overnight.
3. Grill or broil.

## NOTES

Use 8 tablespoons of marinade for every 1 pound of meat. This recipe is enough for 2 pounds.



# Roasted Garlic

Servings: 36

Serving size: 1 clove



LOW  
SODIUM



LOW  
PROTEIN



REDUCES  
ACID

## INGREDIENTS

3 heads of garlic

water

## COOKING INSTRUCTIONS

1. Preheat oven to 350° F.
2. Remove skin from garlic heads, leaving cloves intact.
3. Cut top off each head. Put heads in a small baking dish and add water to cover the bottom (about ¼ inch).
4. Cover with a lid or foil.
5. Bake for 60 minutes or until cloves are soft to the touch.

## SERVING SUGGESTION

To eat, squeeze the large end of the clove until it pops out. Use as a spread on bread, substitute for mayonnaise on a sandwich, or as a topping for pizza or baked potato. Leftover cloves will keep in a sealed container in the refrigerator for about a week.

### NUTRITION INFORMATION

(per serving)

Calories 5

Fat 0 g

Carbohydrates 1 g

Protein 0 g

Dietary Fiber 0.1 g

Calcium 5 mg

Phosphorus 5 mg

Sodium 1 mg

Potassium 12 mg





LOW  
SODIUM



LOW  
PROTEIN



HIGH  
FLUID

# Basic Chicken Stock

Servings: 10

Serving size: 1 cup

Recipe submitted by Geno Leage, PKD chapter volunteer

## INGREDIENTS

- |   |  |
|---|--|
| 1 pound chicken bones,<br>washed, roasted   | 2 stalks celery, washed,<br>coarsely chopped |
| 3 tablespoons olive oil<br>(divided)        | 2 cloves garlic, peeled                      |
| 1 yellow onion, peeled,<br>coarsely chopped | 2 bay leaves                                 |
| 2 carrots, peeled, coarsely<br>chopped      | 1 tablespoon black<br>peppercorns            |
|   | 2 ½ quarts water                             |

### NUTRITION INFORMATION

(per serving)

Calories 65

Fat 4 g

Carbohydrates 3 g

Protein 4 g

Dietary Fiber 0 g

Calcium 20 mg

Phosphorus 27 mg

Sodium 32 mg

Potassium 102 mg

*Nutrient information  
may not be exact  
since transfer of  
nutrients from  
food to the stock is  
estimated.*

## COOKING INSTRUCTIONS

1. Preheat oven to 400° F.
2. Wash the chicken bones and place them in a bowl with 2 tablespoons olive oil to coat the bones.
3. Place the bones on a sheet tray or suitable roasting pan and place in the oven for 30 minutes or until golden brown.
4. Place a gallon-size stock pot on the stove top with 1 tablespoon olive oil and heat on high.
5. Add onion, carrots, celery, garlic, bay leaves and peppercorns to the stock pot and sauté for 10 minutes.
6. Add the bones and cover with water.
7. Bring to a boil and reduce heat to a simmer.
8. Simmer the stock for 45 minutes and then strain into a heat-resistant container.

## SERVING SUGGESTION

When cooled, you can store the chicken stock for 1 week in the refrigerator or up to 6 months in the freezer.



# Glossary of Terms and Topics

**Animal-source protein:** Chicken, pork, beef, fish, eggs and cheese/milk. This also includes less commonly eaten meats such as venison, boar, duck, goose, moose and bison.

**ADPKD:** Autosomal dominant polycystic kidney disease, one of the most common life-threatening genetic diseases, causes fluid-filled cysts to develop and enlarge in both kidneys.

**ARPKD:** Autosomal recessive polycystic kidney disease affects approximately 1 in 20,000 children and can cause death in the first month of life.

**Caffeine:** Caffeine has been shown to have the potential to accelerate cyst growth in ADPKD.<sup>9</sup> However, no formal recommendation has been made related to how much caffeine needs to be restricted or whether patients should avoid it. Recently, a study of 102 ADPKD patients showed no association between low intake of caffeine (average caffeine intake of 86 milligrams per day) with renal volume.<sup>10</sup> Because caffeine is known to promote the growth of cysts, it should be limited in individuals living with ADPKD. However, the consumption of a single cup of coffee (eight ounces equals 70 to 200 milligrams), two cups of tea (16 ounces equals 30 to 140 milligrams) or a soda (20 ounces equals 58 to 100 milligrams) daily may be acceptable. Decaf or caffeine-free versions of drinks are more appropriate. If possible, they should be selected instead of caffeinated beverages. Caffeine-containing supplements, energy shots and energy drinks should be avoided.

**Calcium:** Getting your recommended intake of calcium daily will help you maintain strong bones. Calcium can be found in most dairy products.



**Dietary acid load:** A measure of the amount of protein-to-potassium an individual eats. Eating too much protein and not enough potassium-rich foods will result in a high dietary acid load.

**Dietary supplements, including Chinese herbs:** Many individuals may take supplements in addition to their medications. **Supplements can potentially be dangerous. Unlike drugs, dietary supplements are treated like special foods and are not regulated by the same guidelines.** One key difference is that drugs are considered unsafe until proven safe through rigorous scientific trials, whereas dietary supplements are considered safe by default and must be proven unsafe. This means that no safety or efficacy trials are needed before marketing a supplement. Supplements may or may not contain the ingredients that are listed on the label. A 2013 study that sampled 44 herbal supplements found that only 48 percent contained any of the herbs listed on the label.<sup>43</sup> Of the ones that contained the herb, most had fillers or were contaminated with other products. Many supplements interact with medications that may reduce the drug's effectiveness or cause unknown side effects. Before taking a supplement, talk to your doctor about what you are considering. Fish oil, multivitamins and calcium supplements are often the only supplements prescribed by physicians, typically on an individual basis.

**Dietitian versus nutritionist:** Eating healthy is important for individuals with PKD, but when it comes to diet and nutrition counseling, from whom should you be getting your information? Many people hear the terms "dietitian" and "nutritionist" and think they are the same; however, there are some important differences. To clear up some of the confusion, the Academy of Nutrition and Dietetics, the world's largest organization of food and dietetics practitioners, has stated that **"all registered dietitians are nutritionists but not all nutritionists are registered dietitians."**<sup>44</sup> Unlike nutritionists, dietitians must adhere to strict education and training requirements set forth by a governing body, making them the only nutrition professionals regulated by law. This is why dietitians are the only nutrition experts legally able to provide medical nutrition therapy to patients.





To become a dietitian, individuals must obtain a certain set of qualifications:

- Complete a bachelor's degree at a university accredited or approved by the Accreditation Council for Education in Nutrition and Dietetics. Dietitians will require a master's degree as of 2024.
- Complete 1,200 hours of supervised practice in a variety of clinical, community and food-service rotations.
- Pass a national examination administered by the Commission on Dietetic Registration.
- Complete ongoing continuing professional education that is reviewed by the Commission on Dietetic Registration over the course of their entire career.

Dietitians will have either the registered dietitian (RD) or registered dietitian nutritionist (RDN) credential after their names. They both indicate the same credential and qualifications.

In most states, the term "nutritionist" isn't protected, meaning that no formal requirements exist to use the credential. This means that someone with no formal education, training or expertise in nutrition can call him or herself a nutritionist. A few states have started protecting the term "nutritionist" to avoid issues with patients getting advice from unqualified individuals; however, only a handful of states have been able to do so.

Just like doctors, dietitians often have a particular field of expertise, such as cancer, kidney disease or diabetes. When looking for a dietitian, it is better to find one who specializes in the particular condition that you have. With PKD, it is best if you can find a dietitian who specializes in nephrology; however, if you live in a rural area, this may not always be possible.

Your insurance may not cover seeing an RD or RDN about your dietary needs. Check with your insurance company first to see what services may be covered and the best route to get them covered. Even if services aren't covered, you should consider seeing an RD if you want to make dietary changes. He or she will help you figure out what is important to you and your health, determine how best to approach making these changes and set up a plan based on your medical needs and the most current evidence-based recommendations. The RD also



can help you follow through on the plan. Although you may have to pay out of pocket, changing your dietary behaviors may help you stay out of the hospital and save you money in future medical expenses, making it an investment in your long-term health.

When looking for advice on diet, check whether the individual has the RD or RDN credential after his or her name. These credentials indicate that the individual has completed the required education and training to give nutrition advice to patients. These are the only legally accepted credentials for health care professionals who specialize in nutrition. You can find an expert to help you with your diet by going to [eatright.org/find-an-expert](http://eatright.org/find-an-expert) and searching by ZIP code and specialty.

**HDL cholesterol:** High-density lipoprotein cholesterol. Also known as the “good” cholesterol.

**KDOQI guidelines:** Kidney Disease Outcomes Quality Initiative guidelines are evidence-based clinical practice guidelines for all stages of chronic kidney disease.

**nPCR level:** The normalized protein catabolic rate level measures if you have enough protein in your diet; used in hemodialysis.

**nPNA level:** The normalized protein nitrogen appearance level measures if you have enough protein in your diet; used in peritoneal dialysis.

**Nutritionist:** An individual who advises on matters of food and diet. A nutritionist does not necessarily hold a professional credential or have formal training. See page 110 for more information.

**Phosphorus:** A mineral found naturally in many foods, primarily dairy products and meat. Phosphorus is also abundant in processed and packaged foods, since it acts as a preservative and can increase a food's shelf-life. The kidneys play an important role in keeping phosphorus levels in the blood in normal range. When kidney function declines, you may need to limit your intake of foods high in



phosphorus. Your physician may prescribe phosphorus binders to be taken with meals as well (see phosphorus binders).

**Phosphorus binders:** These help prevent phosphorus from the foods that you eat from being absorbed.

**Potassium:** May play an important role in protecting the kidneys by reducing your dietary acid load. It is an important nutrient for keeping your blood pressure in normal range. Potassium may need to be restricted as kidney function declines.

**Registered Dietitian (RD):** A food and nutrition expert who has met rigorous qualifying requirements. See page 110 for more information.

**Registered Dietitian Nutritionist (RDN):** See *Registered Dietitian*.

**Serum phosphorus levels:** A measurement of phosphorus in the blood.

**Vasopressin:** A naturally occurring hormone that functions to properly regulate fluid balance in our bodies. It helps to conserve water when we are dehydrated and excrete water when we have too much. High vasopressin levels, which occur when we don't get enough fluid, have been linked to many diseases, including high blood pressure, heart and kidney disease. High vasopressin levels have a particularly harmful effect in ADPKD.

**Water loading:** Water has gained significant interest as a therapy for individuals with PKD. Drinking water reduces levels of vasopressin, a hormone that stimulates cyst growth.<sup>6</sup> Vasopressin is stimulated during times of dehydration or after a high-sodium or high-protein meal.<sup>7</sup> Although drinking a large volume of water once or twice a day may lower your vasopressin levels in the short term for individuals living with PKD, **water should be consumed generously throughout the day to keep vasopressin levels consistently low.** For adults with ADPKD, a minimum of three liters (100 ounces) of water should be consumed daily.<sup>5</sup>



# References

1. Torres, V.E., Grantham, J.J., Chapman, A.B., et al. "Potentially Modifiable Factors Affecting the Progression of Autosomal Dominant Polycystic Kidney Disease," *Clinical Journal of the American Society of Nephrology* Mar 2011, 6(3): 640-647.
2. Cowley, B.D., Jr., Grantham, J.J., Muessel, M.J., Kraybill, A.L., Gattone, V.H., 2nd. "Modification of Disease Progression in Rats With Inherited Polycystic Kidney Disease," *American Journal of Kidney Diseases* Jun 1996, 27(6): 865-879.
3. Tanner, G.A., Tanner, J.A. "Citrates Therapy for Polycystic Kidney Disease in Rats," *Kidney International* Nov 2000, 58(5): 1859-1869.
4. Nagao, S., Nishii, K., Katsuyama, M., et al. "Increased Water Intake Decreases Progression of Polycystic Kidney Disease in the PCK Rat," *Journal of the American Society of Nephrology* Aug 2006, 17(8): 2220-2227.
5. Wang, C.J., Creed, C., Winklhofer, F.T., Grantham, J.J. "Water Prescription in Autosomal Dominant Polycystic Kidney Disease: A Pilot Study," *Clinical Journal of the American Society of Nephrology* Jan 2011, 6(1): 192-197.
6. Torres, V.E., Harris, P.C. "Strategies Targeting cAMP Signaling in the Treatment of Polycystic Kidney Disease," *Journal of the American Society of Nephrology* Jan 2014, 25(1): 18-32.
7. Torres, V.E., Bankir, L., Grantham, J.J. "A Case for Water in the Treatment of Polycystic Kidney Disease," *Clinical Journal of the American Society of Nephrology* Jun 2009, 4(6): 1140-1150.
8. Ogborn, M.R., Sareen, S. "Amelioration of Polycystic Kidney Disease by Modification of Dietary Protein Intake in the Rat," *Journal of the American Society of Nephrology* Dec 1995, 6(6): 1649-1654.
9. Belibi, F.A., Wallace, D.P., Yamaguchi, T., Christensen, M., Reif, G., Grantham, J.J. "The Effect of Caffeine on Renal Epithelial Cells From Patients With Autosomal Dominant Polycystic Kidney Disease," *Journal of the American Society of Nephrology* Nov 2002, 13(11): 2723-2729.
10. Vendramini, L.C., Nishiura, J.L., Baxmann, A.C., Heilberg, I.P. "Caffeine Intake by Patients With Autosomal Dominant Polycystic Kidney Disease," *Brazilian Journal of Medical and Biological Research* Sep 2012, 45(9): 834-840.
11. Oldrizzi, L., Rugiu, C., Valvo, E., et al. "Progression of Renal Failure in Patients With Renal Disease of Diverse Etiology on Protein-Restricted Diet," *Kidney International* Mar 1985, 27(3): 553-557.
12. Goraya, N., Simoni, J., Jo, C., Wesson, D.E. "Dietary Acid Reduction With Fruits and Vegetables or Bicarbonate Attenuates Kidney Injury in Patients With a Moderately Reduced Glomerular Filtration Rate Due to Hypertensive Nephropathy," *Kidney International* Jan 2012, 81(1): 86-93.
13. Phisitkul, S., Khanna, A., Simoni, J., et al. "Amelioration of Metabolic Acidosis in Patients With Low GFR Reduced Kidney Endothelin Production and Kidney Injury, and Better Preserved GFR," *Kidney International*. Apr 2010, 77(7): 617-623.
14. Mahajan, A., Simoni, J., Sheather, S.J., Broglio, K.R., Rajab, M.H., Wesson, D.E. "Daily Oral Sodium Bicarbonate Preserves Glomerular Filtration Rate by Slowing Its Decline in Early Hypertensive Nephropathy," *Kidney International* Aug 2010, 78(3): 303-309.



15. Banerjee, T., Crews, D.C., Wesson, D.E., et al. "High Dietary Acid Load Predicts ESRD Among Adults With CKD," *Journal of the American Society of Nephrology* Feb 12 2015, 10: 1681.
16. USRDS 2014 Annual Data Report: "Atlas of Chronic Kidney Disease and End-Stage Renal Disease in the United States," 2014.
17. Szinnai G, Morgenthaler NG, Berneis K, Struck J, Muller B, Keller U, Christ-Crain M. "Changes in Plasma Copeptin, the C-terminal Portion of Arginine Vasopressin During Water Deprivation and Excess in Healthy Subjects," *J Clin Endocrinol Metab* 92: 3973–3978, 2007
18. American Heart Association. "Shaking the Salt Habit," 2014; [http://www.heart.org/HEARTORG/Conditions/HighBloodPressure/PreventionTreatmentofHighBloodPressure/Shaking-the-Salt-Habit\\_UCM\\_303241\\_Article.jsp](http://www.heart.org/HEARTORG/Conditions/HighBloodPressure/PreventionTreatmentofHighBloodPressure/Shaking-the-Salt-Habit_UCM_303241_Article.jsp). Feb 2014.
19. Frassetto, L.A., Todd, K.M., Morris, R.C., Jr., Sebastian, A. "Estimation of Net Endogenous Noncarbonic Acid Production in Humans From Diet Potassium and Protein Contents," *The American Journal of Clinical Nutrition* Sep 1998, 68(3): 576-583.
20. Appel, L.J., Moore, T.J., Obarzanek, E., et al. "A Clinical Trial of the Effects of Dietary Patterns on Blood Pressure, DASH Collaborative Research Group," *The New England Journal of Medicine* Apr 17 1997, 336(16): 1117-1124.
21. "Dietary Reference Intakes (DRIs): Recommended Dietary Allowances and Adequate Intakes, Vitamins," 2012, <http://www.iom.edu/Global/News%20Announcements/-/media/C5CD2DD7840544979A549EC47E56A02B.ashx>.
22. "Dietary Reference Intakes for Energy, Carbohydrates, Fiber, Fat, Fatty Acids, Cholesterol, Protein and Amino Acids," 2002/2005, <http://www.iom.edu/Global/News%20Announcements/-/media/C5CD2DD7840544979A549EC47E56A02B.ashx>.
23. Kokkinos, P.F., Fernhall, B. "Physical Activity and High Density Lipoprotein Cholesterol Levels: What Is the Relationship?" *Sports Medicine* Nov 1999, 28(5): 307-314.
24. Craig, W.Y., Palomaki, G.E., Haddow, J.E. "Cigarette Smoking and Serum Lipid and Lipoprotein Concentrations: An Analysis of Published Data. *BMJ*. Mar 25, 1989, 298(6676): 784-788.
25. Verbalis J. "Disorders of Water Balance" In: Saunders, ed. *Brenner and Rector's The Kidney* 9th ed, Elsevier, Inc., 2012.
26. Amro, O.A., Paulus, J.K., Noubary, F., Perrone, R.D. "Low Osmolar Diet and Adjusted Water Intake for Vasopressin Suppression in ADPKD: A Randomized Controlled Trial," *ASN Kidney Week* 2015, Nov 3-8, 2015; San Diego, CA.
27. Kahn, H.D., Stralka, K. "Estimated Daily Average Per Capita Water Ingestion by Child and Adult Age Categories Based on USDA's 1994-1996 and 1998 Continuing Survey of Food Intakes by Individuals," *Journal of Exposure Science and Environmental Epidemiology*. May 2009, 19(4): 396-404.
28. Taylor, J.M., Hamilton-Reeves, J.M., Sullivan, D.K., et al. "Diet and Polycystic Kidney Disease: A Pilot Intervention Study," *Clinical Nutrition* Jan 11, 2016.
29. Tantamango-Bartley, Y., Jaceldo-Siegl, K., Fan, J., Fraser, G. "Vegetarian Diets and the Incidence of Cancer in a Low-Risk Population," *Cancer Epidemiology, Biomarkers and Prevention*, a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology, Feb 2013, 22(2):286-294.
30. Szeto, Y.T., Kwok, T.C., Benzie, I.F. "Effects of a Long-Term Vegetarian Diet on Biomarkers of Antioxidant Status and Cardiovascular Disease Risk," *Nutrition* Oct 2004;20(10):863-866.



31. Sabate, J., Wien, M. "Vegetarian Diets and Childhood Obesity Prevention," *The American Journal of Clinical Nutrition* May 2010, 91(5): 1525S-1529S.
32. Craig, W.J., Mangels, A.R. "American Dietetic Association. Position of the American Dietetic Association: Vegetarian Diets" *Journal of the American Dietetic Association* Jul 2009, 109(7): 1266-1282.
33. Katcher, H.I., Ferdowsian, H.R., Hoover, V.J., Cohen, J.L., Barnard, N.D. "A Worksite Vegan Nutrition Program Is Well-Accepted and Improves Health-Related Quality of Life and Work Productivity," *Annals of Nutrition and Metabolism*. 2010, 56(4): 245-252.
34. Rastegar, A. "Serum Potassium. In: Walker, H.K., Hall, W.D., Hurst, J.W. "Clinical Methods: The History, Physical and Laboratory Examinations," 3rd ed., Boston 1990.
35. National Kidney Foundation. "KDOQI Clinical Practice Guidelines for Bone Metabolism and Disease in Chronic Kidney Disease Guideline 3," "Evaluation of Serum Phosphorus Levels," *American Journal of Kidney Disease* 2003,42: S62-S63.
36. Workgroup KD. "KDOQI Clinical Practice Guidelines for Cardiovascular Disease in Dialysis Patients," *American Journal of Kidney Diseases* Apr 2005, 45 (4 Suppl 3): S1-153.
37. KDOQI Work Group. "KDOQI Clinical Practice Guideline for Nutrition in Chronic Renal Failure," *American Journal of Kidney Disease*. 2000, 35(6): S44-S45.
38. Bethke, P.C., Jansky, S.H. "The Effects of Boiling and Leaching on the Content of Potassium and Other Minerals in Potatoes," *Journal of Food Science* Jun 2008, 73(5): H80-85.
39. Ando, S., Sakuma, M., Morimoto, Y., Arai, H. "The Effect of Various Boiling Conditions on Reduction of Phosphorus and Protein in Meat," *Journal of Renal Nutrition*, the official journal of the Council on Renal Nutrition of the National Kidney Foundation, Nov 2015, 25(6): 504-509.
40. van den Berg, E., Engberink, M.F., Brink, E.J., et al. "Dietary Acid Load and Metabolic Acidosis in Renal Transplant Recipients," *Clinical Journal of the American Society of Nephrology* Nov 2012, 7(11): 1811-1818.
41. Costa, B., Moratelli, L., Silva, L.B., et al. "Body Mass Index in the First Year After Kidney Transplantation," *Transplantation Proceedings* Jul-Aug 2014, 46(6): 1750-1752.
42. Hricik, D.E. "Metabolic Syndrome in Kidney Transplantation: Management of Risk Factors," *Clinical Journal of the American Society of Nephrology* Jul 2011,6(7): 1781-1785.
43. Newmaster, S.G., Grguric, M., Shanmughanandhan, D., Ramalingam, S., Ragupathy, S. "DNA Barcoding Detects Contamination and Substitution in North American Herbal Products," *BMC Medicine*, 2013, 11:222.
44. Academy of Nutrition and Dietetics. "Every Registered Dietitian Is a Nutritionist, but Not Every Nutritionist Is a Registered Dietitian. [www.eatrightpro.org/resource/about-us/what-is-an-rdn-and-dtr/what-is-a-registered-dietitian-nutritionist/every-registered-dietitian-is-a-nutritionist-but-not-every-nutritionist-is-a-registered-dietitian](http://www.eatrightpro.org/resource/about-us/what-is-an-rdn-and-dtr/what-is-a-registered-dietitian-nutritionist/every-registered-dietitian-is-a-nutritionist-but-not-every-nutritionist-is-a-registered-dietitian).
45. Wardle, J., Cooke L.J., Gibson, E.L., Sapochnik, M., Sheiham, A., Lawson, M. "Increasing Children's Acceptance of Vegetables: A Randomized Trial of Parent-Led Exposure," *Appetite* Apr 2003, 40(2): 155-162.
46. Wardle, J., Herrera, M.L., Cooke, L., Gibson, E.L. "Modifying Children's Food Preferences: The Effects of Exposure and Reward on Acceptance of an Unfamiliar Vegetable," *European Journal of Clinical Nutrition* Feb 2003, 57(2): 341-348.



# Index

## **APPLE**

- Apple Muffin 99
- Rosemary Sage Burgers With  
Apple Slaw and Chive Mayo 76
- Spinach, Apple and  
Walnut Salad 59

## **ARTICHOKES**

- Artichoke Dip 34

## **ASPARAGUS**

- Grilled Asparagus With  
Mozzarella 68

## **AVOCADO**

- Quinoa With Black Beans and  
Avocado 71

## **BEEF**

- Beef Chili 51
- Coffee-Ancho Chile-Rubbed  
Flat Iron Steak 86
- Rosemary Sage Burgers With  
Apple Slaw and Chive Mayo 76
- Sweet and Spicy Meatballs 38

## **BLACK BEANS**

- Quinoa With Black Beans and  
Avocado 71
- Sweet Potato Burgers 79

## **BLACKBERRY**

- Blackberry Cobbler 93

## **BODY WEIGHT**

- Nutrition Guidelines for  
Individuals Living With  
ADPKD 7

## **BONE HEALTH**

- Nutrition Guidelines for  
Individuals With Kidney  
Transplants 23

## **BROCCOLI**

- Mushroom and Broccoli Risotto 87

## **BRUSSELS SPROUTS**

- Roasted Brussels Sprouts 67

## **BURGERS/SANDWICHES/ WRAPS**

- Chicken Lettuce Wraps 36
- Chipotle Chicken Salad 58
- Rosemary Sage Burgers With  
Apple Slaw and Chive Mayo 76
- Sweet Potato Burgers 79

## **CAFFEINE**

- Nutrition Guidelines for  
Individuals Living With  
ADPKD 10

## **CAKE**

- Coconut Angel Food Cake 94

## **CALCIUM**

- Nutrition Guidelines for  
Individuals Living With  
ADPKD 10
- Nutrition Guidelines for  
Individuals on Dialysis 21





## **CALORIES**

Nutrition Guidelines for Individuals Living With ADPKD	11
Nutrition Guidelines for Individuals on Dialysis	21

## **CARROTS**

Creamy Carrot Thyme Soup	50
--------------------------	----

## **CHICKEN**

Basic Chicken Stock	108
Chicken and Spanish Rice	83
Chicken Lettuce Wraps	36
Chipotle Chicken Salad	58
Grilled Malt Chicken With Peppers and Onion	84

## **CHICKPEAS/ GARBANZO BEANS**

Couscous and Feta-Stuffed Peppers	73
Hummus	42

## **CHILDREN WITH PKD**

Nutrition Guidelines for Children at Risk of Having ADPKD and Individuals With ARPKD	26
--	----

## **CHILES/PEPPERS**

Grilled Malt Chicken With Peppers and Onion	84
Roasted Vegetables	72

## **CHOCOLATE**

Chocolate Chip Cookies	90
------------------------	----

## **COCONUT**

Coconut Angel Food Cake	94
-------------------------	----

## **CONDIMENTS**

Homemade Mayonnaise	104
---------------------	-----

## **COOKIES**

Chocolate Chip Cookies	90
------------------------	----

## **CORN**

Roasted Corn and Edamame Salad	54
-----------------------------------	----

## **COUSCOUS**

Couscous and Feta-Stuffed Peppers	73
--------------------------------------	----

## **CRAB MEAT**

Hot Crab Dip	39
Shrimp and Crab Gumbo	81

## **DIPS**

Artichoke Dip	34
Fresh Tzatziki	35
Hot Crab Dip	39
Hummus	42
Light and Creamy Cranberry Fresh Fruit Dip	92
Mexican Layer Dip	40

## **EDAMAME**

Roasted Corn and Edamame Salad	54
-----------------------------------	----

## **EGGPLANT**

Eggplant Vegetable Soup	47
-------------------------	----

## **EXERCISE**

Nutrition Guidelines for Individuals Living With ADPKD	11
--	----

## **FLUIDS**

Nutrition Guidelines for Individuals Living With ADPKD	9
Nutrition Guidelines for Individuals on Dialysis	20





## **FOOD SAFETY**

Nutrition Guidelines for  
Individuals With Kidney  
Transplants 24

## **GINGER**

Ginger Pork Soup 49

## **HERBAL SUPPLEMENTS**

Nutrition Guidelines for  
Individuals With Kidney  
Transplants 23

## **KALE**

Warm Potato and Kale Mix 63

## **LEMON**

Lemon Ricotta Pie 96  
Zesty Lemon Tart 101

## **LIME**

Chili Lime Shrimp 44  
Key Lime Pie 100  
Strawberry Lime Dessert 91

## **MANDARINS**

Mandarin Salad 57

## **MANGO**

Mango Salsa Wontons 33

## **MARINADE**

Simple Kebob Marinade 106

## **MUFFINS**

Apple Muffin 99  
Bran Muffin 98

## **MUSHROOMS**

Mushroom and Broccoli Risotto 87

## **PAPAYA**

Grilled Salmon With  
Papaya-Mint Salsa 75

## **PASTA**

Fall Harvest Orzo Salad 61  
Noodles Romano 70

## **PEACH**

Peach Pork Tenderloin 77

## **PHOSPHORUS**

Nutrition Guidelines for  
Individuals on Dialysis 20  
Pies/Tarts  
Key Lime Pie 100  
Lemon Ricotta Pie 96  
Zesty Lemon Tart 101

## **PINEAPPLE**

Salmon With Pineapple Salsa 78

## **PORK**

Ginger Pork Soup 49  
Peach Pork Tenderloin 77  
Rosemary Sage Burgers With  
Apple Slaw and Chive Mayo 76

## **POTASSIUM**

Nutrition Guidelines for  
Individuals Living With  
ADPKD 8  
Nutrition Guidelines for  
Individuals on Dialysis 19

## **POTATOES**

Mashed Tomato Potatoes 65  
Roasted Vegetables 72  
Warm Potato and Kale Mix 63





## **PROTEIN**

Nutrition Guidelines for Individuals Living With ADPKD	8
Nutrition Guidelines for Individuals on Dialysis	18



## **QUINOA**

Quinoa With Black Beans and Avocado	71
--	----

## **RICE**

Chicken and Spanish Rice	83
Mushroom and Broccoli Risotto	87

## **SALMON**

Baked Salmon	80
Grilled Salmon With Papaya-Mint Salsa	75
Salmon With Pineapple Salsa	78

## **SEASONINGS**

Coffee-Ancho Chile Rub	105
Homemade Salt-Free Mexican Spice Mix	103
Roasted Garlic	107

## **SHRIMP**

Chili Lime Shrimp	44
Shrimp and Crab Gumbo	81
Summer Shrimp	85

## **SODIUM**

Nutrition Guidelines for Individuals Living With ADPKD	7
Nutrition Guidelines for Individuals on Dialysis	17



## **SPINACH/LETTUCE**

Mandarin Salad	57
Spinach, Apple and Walnut Salad	59
Strawberry Spinach Salad	53
Summertime Salad	55

## **SQUASH**

Grilled Vegetables	69
Roasted Vegetables	72
Zucchini-Ribbon Salad	56

## **STOCK**

Basic Chicken Stock	108
---------------------	-----

## **STRAWBERRY**

Strawberry Lime Dessert	91
Strawberry Spinach Salad	53
Strawberry-Mango Parfaits With Ginger Topping	89

## **SWEET POTATOES**

Grilled Sweet Potatoes and Scallions	66
Grilled Vegetables	69
Roasted Vegetables	72
Spicy Sweet Potato Fries	64
Sweet Potato Burgers	79

## **TOMATOES**

Mashed Tomato Potatoes	65
Mexican Vegetable Salad	60

## **TURKEY**

Eggplant Vegetable Soup	47
Turkey Vegetable Soup	48

## **ZUCCHINI**

Grilled Vegetables	69
Zucchini-Ribbon Salad	56





People creating new products  
for better health worldwide

## **THE NAME "OTSUKA" TRANSLATES TO "MAJOR MILESTONE."**

And indeed, for over 95 years, Otsuka-People have achieved major milestones in their quest to create new products for better health.

Today, our commitment to developing new products and innovative solutions to address some of healthcare's most complex challenges has never been stronger.

1.800.562.3974  
[www.otsuka-us.com](http://www.otsuka-us.com)

Otsuka America Pharmaceutical, Inc.  
Otsuka PHarmaceutical Development & Commercialization, Inc.  
Otsuka Maryland Medicinal Laboratotires, Inc.



*Cooking Well* provides a helping hand to those with polycystic kidney disease (PKD) who are looking to play an active role in their health. From appetizers to salads to entrees to desserts and more, these delicious everyday recipes and important nutritional guidelines will help those with PKD live their best life.

“Nutrition is essential for everyone, but especially important for those with polycystic kidney disease. *Cooking Well* is a fantastic resource for patients and families wanting to proactively take charge of their health.”

**Terry Watnick, M.D.**  
*PKD Foundation Science Advisory Committee Chair*

“The nutritious and delicious recipes in this wonderful cookbook make it easy for individuals with PKD and their families to eat in an enjoyable and healthy manner. The menu choices are extensive and the preparation is simple. Good eating!”

**Ronald Perrone, M.D.**  
*Tufts Medical Center, Boston, MA*



**PKD FOUNDATION**  
Polycystic Kidney Disease

This cookbook was made possible with the generous support of Otsuka America Pharmaceutical, Inc.

